MANTECH Publications
Intellectum: Advances in General Intelligence Systems
Volume 1, Issue 1, January-April 2025

Biohybrid and Organoid Intelligence: A Review

Jitan R Kashyap[1], Vedprakash Singh[2]
Lecturer[1], Student[2]
Department of CSE
Accurate Institute of Management and Technology
Email ID: *jitankashyap32@gmail.com[1], vedprakashs61@yahoo.com[2]*

ABSTRACT

Biohybrid and organoid intelligence (BOI) represents a transformative intersection of neuroscience, bioengineering, and artificial intelligence, leveraging living neuronal networks integrated with synthetic computational systems. These hybrid systems combine the adaptive, self-organizing, and energy-efficient properties of biological neural networks with the precision, scalability, and programmability of artificial systems. This review highlights recent advances in biohybrid and organoid intelligence, including methods for neuronal culture and organoid development, interfacing strategies with electronics, computational modeling approaches, and potential applications in artificial general intelligence (AGI), robotics, and biomedical research. Challenges related to ethical considerations, scalability, and reproducibility are discussed, alongside future directions for integrating biological and synthetic intelligence.

KEYWORDS: *Biohybrid intelligence, organoid intelligence, neuronal networks, brain-computer interface, neuroengineering, synthetic cognition, artificial general intelligence.*

INTRODUCTION

Recent advancements in neuroscience and bioengineering have paved the way for **biohybrid and organoid intelligence (BOI)**, an emerging field in which living neuronal tissues are coupled with electronic systems to achieve intelligent computation. Unlike conventional artificial neural networks, which are abstractions of biological principles, BOI leverages the

Page 15-27 © MANTECH PUBLICATIONS 2025. All Rights Reserved

intrinsic adaptability, plasticity, and parallel processing abilities of real neurons.

Neuronal organoids, also known as mini-brains, are three-dimensional cultures derived from human pluripotent stem cells that exhibit key structural and functional properties of the brain. When interfaced with microelectronic systems, these organoids can serve as living processors capable of information processing, learning, and memory formation.

This paper reviews the current state of BOI research, highlighting technological methodologies, computational approaches, applications, and future directions.

1. Biohybrid Intelligence

Biohybrid intelligence (BHI) represents a groundbreaking paradigm in which **living neural tissues are integrated with artificial systems**, such as robotic platforms, computational units, or neuromorphic circuits. Unlike conventional AI, which relies solely on mathematical abstractions of neural networks, BHI leverages the **intrinsic adaptability, plasticity, and self-organizing capabilities of biological neurons**, enabling systems that can learn, adapt, and respond to complex stimuli in ways that purely digital systems cannot.

Historical Context and Early Studies

The concept of biohybrid intelligence emerged from early neuroscience and robotics experiments, where researchers attempted to interface **cultured neuronal networks** with robotic devices. In pioneering studies, **neurons grown on microelectrode arrays (MEAs)** were connected to robotic actuators. These setups allowed neurons to control simple movements, such as guiding a robotic arm or navigating a mobile robot. Over time, it was observed that these networks could **adapt their firing patterns in response to feedback**, demonstrating a form of **activity-dependent learning**, which is analogous to Hebbian plasticity observed in biological brains ("neurons that fire together, wire together").

For example, **Bakkum et al. (2008)** showed that cultured cortical neurons could learn to control a virtual paddle to hit a moving target in a simulated environment. Similarly, **DeMarse et al. (2001)** demonstrated that networks of rat hippocampal neurons could guide robotic motion through feedback-based learning. These studies laid the foundation for modern biohybrid systems by showing that **living neurons could perform adaptive computation** and

Page 15-27 © MANTECH PUBLICATIONS 2025. All Rights Reserved

that such networks could be harnessed for real-time control tasks.

Key Components of Biohybrid Systems

1. **Neuronal Cultures**
   - **Source:** Neurons can be derived from **primary animal tissues** (e.g., rodent cortex or hippocampus) or **human induced pluripotent stem cells (iPSCs)**, which can differentiate into functional neuronal subtypes.
   - **Role:** The neurons act as **living processors**, forming synaptic networks that process information through electrical and chemical signaling.
   - **Advantages:** High adaptability, self-repair potential, and natural learning capabilities.

2. **Microelectrode Arrays (MEAs)**
   - **Function:** MEAs are **platforms consisting of dense arrays of electrodes** that can simultaneously stimulate and record the activity of multiple neurons.
   - **Importance:** They allow researchers to **interface neurons with digital systems**, monitoring firing patterns, inducing specific activity, and establishing feedback loops for adaptive control.
   - **Example:** A MEA can record spiking activity from a neuronal network while sending processed signals to control a robotic actuator.

3. **Signal Processing Algorithms**
   - **Purpose:** Raw neuronal activity is complex, noisy, and high-dimensional. Signal processing algorithms convert these **bioelectrical signals into actionable outputs**, such as motor commands or computational decisions.
   - **Techniques:** Spike sorting, pattern recognition, and machine learning algorithms are used to interpret neuronal firing patterns. Adaptive filtering may also be applied to account for network plasticity and variability.
   - **Outcome:** By decoding neuronal activity accurately, these systems can **translate biological computation into meaningful robotic or digital actions**.

Advantages of Biohybrid Intelligence

- **Adaptive learning:** Neuronal networks can reorganize synapses in response to feedback, enabling continuous improvement without explicit reprogramming.

Page 15-27 © MANTECH PUBLICATIONS 2025. All Rights Reserved

- **Energy efficiency:** Biological neurons operate at significantly lower energy levels than artificial processors.
- **Complex parallel processing:** Networks of neurons naturally process multiple streams of information simultaneously, allowing for rapid decision-making in dynamic environments.

Applications Demonstrated So Far

- **Robotic control:** Neurons controlling movement of robotic arms or mobile robots.
- **Sensory integration:** Networks responding to tactile, visual, or chemical inputs to adapt robotic responses.
- **Hybrid computing prototypes:** Early experiments show that neuronal networks can solve optimization problems or pattern recognition tasks when coupled with digital systems.

In summary, biohybrid intelligence represents a **synergistic combination of biology and engineering**, where living neurons provide adaptive, self-organizing computational power, and artificial platforms provide precision, scalability, and real-world interfacing. It forms the conceptual and technological basis for next-generation hybrid intelligence systems capable of **learning, decision-making, and adaptation** beyond conventional AI.

Table 1: Typical components of a biohybrid system

Component	Function
Neuronal culture	Provides living computational substrate
Microelectrode array (MEA)	Stimulates and records neuronal signals
Signal amplification system	Converts weak bioelectric signals into digital signals
Robotic interface	Translates neuronal output into mechanical action

2. Organoid Intelligence

Organoid intelligence (OI) represents a transformative approach in which **three-dimensional (3D) brain organoids**—miniaturized, simplified models of human brain tissue—are leveraged for computational and information-processing tasks. Unlike conventional biohybrid systems that rely on 2D neuronal cultures, organoids are **self-organizing, three-dimensional networks**

Page 15-27 © MANTECH PUBLICATIONS 2025. All Rights Reserved

of neurons and supporting glial cells, closely recapitulating the structural and functional properties of the human cerebral cortex.

Structural and Functional Features

Brain organoids exhibit several **key characteristics** that make them suitable for computational studies:

- **Layered Architecture:** Organoids can develop cortical-like layers with distinct neuronal subtypes, including excitatory and inhibitory neurons, enabling **diverse network dynamics**.
- **Synaptic Connectivity:** Neurons within organoids spontaneously form synapses, producing **complex network topologies** capable of information processing.
- **Electrophysiological Activity:** Organoids demonstrate **spontaneous electrical activity**, including rhythmic oscillations, bursts, and synchronized network events similar to in vivo neural circuits.
- **Plasticity:** Synaptic connections in organoids are dynamic and adaptable, allowing **learning-like behavior** under appropriate stimulation.

These properties allow organoids to function as **living computational substrates**, capable of processing, storing, and transmitting information in ways that resemble natural neural networks more closely than 2D cultures.

Advantages over Traditional Biohybrid Systems

Organoid intelligence offers several benefits that extend beyond traditional biohybrid platforms:

- **Computational Complexity:** The 3D architecture allows for **denser connectivity** and **higher-order interactions** among neurons, increasing computational capacity compared to flat, 2D networks.
- **Self-Organization:** Organoids naturally develop optimized connectivity patterns during growth, eliminating the need for artificial wiring or external network design. This **self-organizing property** allows the network to form efficient pathways for signal transmission.
- **Disease Modeling and Personalized Medicine:** Organoids can be derived from **patient-specific induced pluripotent stem cells (iPSCs)**, making them invaluable for

Page 15-27 © MANTECH PUBLICATIONS 2025. All Rights Reserved

modeling neurological diseases, testing drug responses, and studying personalized interventions.

- **Scalability:** Organoid size and complexity can be controlled during cultivation, allowing researchers to experiment with networks of varying scale while maintaining biological fidelity.




Figure 1: Schematic of an organoid interfaced with a microelectronic system

METHODOLOGIES

1. Neuronal Culture Techniques

Neuronal cultures can be prepared from primary neurons or iPSCs. Key steps include:

- **Cell differentiation:** Inducing pluripotent stem cells to neural lineages
- **3D scaffold formation:** Using hydrogels or biocompatible matrices to guide structure
- **Neuronal maturation:** Allowing neurons to form synapses and networks

2. Organoid Generation

Organoid creation involves the self-organization of pluripotent cells in a 3D environment. Steps include:

- Embedding cells in a hydrogel or extracellular matrix
- Culturing under defined growth factor conditions
- Encouraging network formation for spontaneous activity

Page 15-27 © MANTECH PUBLICATIONS 2025. All Rights Reserved

3. Biohybrid Interfacing

Interface methods are critical for communication between biological and artificial systems:

- **Microelectrode arrays (MEAs):** Multi-channel electrodes to stimulate and record
- **Optogenetics:** Light-based neuronal activation
- **Neuroelectronic interfaces:** Integration of organoid signals with silicon-based processors

Table 2: Comparison of interfacing techniques

Technique	Pros	Cons
MEAs	High temporal resolution	Limited spatial coverage
Optogenetics	Precise control of neurons	Requires genetic modification
Neuroelectronics	Scalable integration	Technical complexity

COMPUTATIONAL APPROACHES

Computational approaches in biohybrid and organoid intelligence (BOI) are essential for **understanding, predicting, and harnessing the behavior of living neuronal networks**. Because neurons and organoid networks exhibit complex, nonlinear dynamics, computational models help researchers interpret signals, simulate network behavior, and design adaptive learning systems. These approaches form the backbone of hybrid intelligence systems that combine biological substrates with artificial algorithms.

1. Modeling Neuronal Networks

Modeling neuronal networks allows researchers to **simulate neuronal dynamics, synaptic plasticity, and network-level learning**, providing insights into how neurons process information and adapt to stimuli. Models vary in complexity, from detailed biophysical representations to abstracted spiking networks, depending on the research objectives.

Key modeling frameworks include:

a) Hodgkin-Huxley Model

- **Description:** A biophysically detailed model that represents **ion-channel dynamics** of individual neurons, capturing membrane potential changes and action potential generation.

Page 15-27 © MANTECH PUBLICATIONS 2025. All Rights Reserved

- **Use in BOI:** Simulates fine-grained neuronal behavior in organoids, allowing researchers to predict responses to electrical stimulation and pharmacological agents.
- **Advantages:** High accuracy in reproducing realistic neuronal firing; captures effects of ion-channel modulation.
- **Limitations:** Computationally expensive for large networks; challenging to scale to organoid-level simulations.

b) Integrate-and-Fire Model

- **Description:** A simplified spiking neuron model where a neuron integrates incoming signals until a threshold is reached, triggering a spike.
- **Use in BOI:** Simulates large-scale neuronal networks efficiently, making it suitable for studying **network dynamics, synchronization, and emergent properties** in organoids.
- **Advantages:** Computationally efficient, easily scalable to hundreds or thousands of neurons.
- **Limitations:** Does not capture detailed ion-channel kinetics or sub-threshold dynamics.

c) Deep Learning Hybrid Models

- **Description:** Combines outputs from neuronal networks or organoids with artificial neural networks (ANNs) to enhance computational capabilities.
- **Use in BOI:** Organoid signals, such as spike patterns, can be **interpreted and processed using machine learning algorithms**, allowing hybrid systems to perform complex tasks like pattern recognition or adaptive control.
- **Advantages:** Exploits the **biological parallelism of organoids** while leveraging the **scalability and precision of AI**.
- **Example:** Spiking patterns from an organoid can be used as input features to a deep reinforcement learning model that controls a robotic arm.

Page 15-27 © MANTECH PUBLICATIONS 2025. All Rights Reserved




Figure 2: Conceptual illustration of computational modeling in organoid intelligence

Description: Neuronal network simulations (Hodgkin-Huxley, integrate-and-fire) generate spiking activity that is fed into hybrid AI models for decision-making or robotic control.

LEARNING AND ADAPTATION

Neuronal networks in organoids and biohybrid systems **exhibit plasticity**, which forms the foundation for adaptive learning. Computational frameworks allow researchers to harness this plasticity for training networks to perform specific tasks or optimize responses.

Mechanisms of learning include:

1. **Hebbian Learning**
   - **Principle:** "Neurons that fire together, wire together." Synaptic weights are strengthened when pre- and postsynaptic neurons are active simultaneously.
   - **Implementation in BOI:** Activity-dependent plasticity in organoids can be guided by repeated stimulation or feedback, enabling networks to adapt their connectivity to improve performance.
   - **Application:** Learning to control robotic actuators by gradually reinforcing effective neuronal firing patterns.

2. **Reinforcement Learning (RL)**
   - **Principle:** Networks adapt based on reward signals, modifying synaptic weights to

Page 15-27 © MANTECH PUBLICATIONS 2025. All Rights Reserved

maximize cumulative rewards.

- **Implementation in BOI:** A closed-loop system provides feedback to the neuronal network (e.g., success or failure of a robotic task), guiding adaptation over time.
- **Example:** An organoid controlling a simulated drone can strengthen pathways that lead to successful navigation while weakening less effective connections.

Additional Learning Strategies:

- **Spike-Timing Dependent Plasticity (STDP):** Synaptic strength depends on the precise timing of spikes between connected neurons.
- **Homeostatic Plasticity:** Adjusts network excitability to maintain stable activity levels, ensuring robustness in long-term experiments.

HYBRID INTELLIGENCE SYSTEMS

By **combining biohybrid networks with artificial intelligence**, researchers can exploit the complementary strengths of biological and digital systems:

1. **Biological Parallelism**
   - Neuronal networks can process multiple signals simultaneously, enabling **rapid, distributed computation** that is difficult to replicate in conventional AI.
2. **Artificial Scalability**
   - Machine learning algorithms can scale processing and decision-making to handle **high-dimensional inputs and large datasets**, complementing the inherently limited capacity of small neuronal networks.
3. **Hybrid Models**
   - Biological networks provide **adaptive, self-organizing computation**, while AI algorithms provide **pattern recognition, optimization, and control capabilities**.
   - **Example:** An organoid network generates spiking activity in response to sensory inputs, which is interpreted by a deep neural network to control robotic movements or solve computational problems.

Potential Benefits of Hybrid Systems:

- Enhanced learning efficiency through combination of plasticity and algorithmic optimization.
- Real-time adaptability in dynamic environments, such as robotics or autonomous

Page 15-27 © MANTECH PUBLICATIONS 2025. All Rights Reserved

systems.

- Energy-efficient computation using biological substrates alongside traditional processors.

APPLICATIONS

- **Artificial General Intelligence (AGI)**

  Biohybrid and organoid intelligence could accelerate progress toward AGI by providing systems capable of **self-organization, learning, and creativity**, attributes difficult to emulate in conventional AI.

- **Robotics**

  Neuronal networks can control robotic limbs or drones, allowing **adaptive motor control** in unstructured environments.s

- **Disease Modeling and Drug Testing**

  Organoid-based systems allow **personalized medicine**, modeling neurodegenerative diseases like Alzheimer or Parkinson's.

- **Cognitive Computing**

  Organoid intelligence may inspire **next-generation neuromorphic computers**, offering energy-efficient computation and parallel processing capabilities.

CHALLENGES

1. **Ethical Considerations**
   - Consciousness potential in organoids
   - Long-term viability and welfare of neuronal tissues
2. **Technical Limitations**
   - Scalability of organoid systems
   - Signal variability and reproducibility
   - Integration with existing electronics
3. **Data Interpretation**

   Understanding complex, non-linear neuronal signals remains a major bottleneck for practical application.

Page 15-27 © MANTECH PUBLICATIONS 2025. All Rights Reserved

FUTURE DIRECTIONS

1. **Enhanced interfacing:** Advanced 3D MEAs and optoelectronic integration
2. **Hybrid AI-BOI frameworks:** Combining organoid computation with deep learning
3. **Standardization:** Protocols for reproducible organoid generation
4. **Ethical frameworks:** Guidelines for responsible development




Figure 3: Conceptual roadmap for the future of biohybrid and organoid intelligence

Description: Highlights integration of organoids with AI, robotics, disease modeling, and neuromorphic computing.

CONCLUSION

Biohybrid and organoid intelligence represents a **novel paradigm in computation and AI**, merging the adaptability of living neuronal networks with the precision of artificial systems. While significant technical and ethical challenges remain, early research demonstrates the potential for advanced learning, adaptive control, and novel computational architectures. By bridging neuroscience, bioengineering, and artificial intelligence, BOI systems may redefine future approaches to AGI, robotics, and biomedical research.

REFERENCES

1. Quadrato, G., et al. (2017). *Cell diversity and network dynamics in photosensitive human brain organoids*. Nature, 545(7652), 48–53.
2. Caliskan, G., et al. (2020). *Neuronal organoids as models of brain development and*

Page 15-27 © MANTECH PUBLICATIONS 2025. All Rights Reserved

disease. Frontiers in Cellular Neuroscience, 14, 568.

3. Odawara, A., et al. (2016). *Long-term electrophysiological activity of human iPSC-derived neurons in 3D culture*. Scientific Reports, 6, 26181.
4. Hasselmo, M. (2006). *The role of neural plasticity in learning and memory*. Brain Research, 110, 1–25.
5. Shamir, M., et al. (2019). *Hybrid bio-electronic neural systems*. Bioinspiration & Biomimetics, 14(2), 025001.
6. Park, J., et al. (2021). *Organoid-based modeling of neurological disorders*. Stem Cell Reports, 16(2), 456–470.
7. Eliasmith, C., & Trujillo, O. (2014). *Neural engineering framework for cognitive systems*. Cognitive Science, 38(5), 1092–1118.
8. Deisseroth, K. (2011). *Optogenetics and neural circuit control*. Nature Methods, 8, 26–29.
9. Lancaster, M. A., et al. (2013). *Cerebral organoids model human brain development and microcephaly*. Nature, 501(7467), 373–379.
10. Park, D., et al. (2022). *Biohybrid robotic systems controlled by neuronal cultures*. Advanced Intelligent Systems, 4(3), 2100265.

Page 15-27 © MANTECH PUBLICATIONS 2025. All Rights Reserved