

Phytochemical Investigation and Pharmacognostic Evaluation of Selected Medicinal Plants in Nigeria

Dr. Chinedu Okafor¹, Amina Bello², Ibrahim Musa³

Associate Professor¹, Students^{2, 3}

Department of Pharmacognosy

University of Nigeria, Nsukka

Corresponding Author Email: chineduokafor@yahoo.com¹

DOI: <https://doi.org/10.5281/zenodo.19707600>

ABSTRACT

*Nigeria possesses a vast diversity of medicinal plants widely used in traditional healthcare systems. Pharmacognosy and phytochemical sciences play a crucial role in validating these indigenous remedies through scientific evaluation. This study focuses on the pharmacognostic standardization and phytochemical screening of selected Nigerian medicinal plants including *Azadirachta indica*, *Vernonia amygdalina*, and *Ocimum gratissimum*. Standard procedures such as macroscopic, microscopic, and physicochemical analyses were performed. Phytochemical investigations revealed the presence of alkaloids, flavonoids, tannins, saponins, and glycosides, indicating significant therapeutic potential. The study highlights the importance of integrating traditional knowledge with modern analytical techniques to ensure drug safety, efficacy, and quality control.*

KEYWORDS: *Pharmacognosy, Phytochemistry, Nigerian medicinal plants, Herbal medicine, Bioactive compounds*

INTRODUCTION

Nigeria, located in West Africa, is endowed with a rich biodiversity comprising thousands of plant species, many of which possess medicinal value. The reliance on herbal medicine in Nigeria is deeply rooted in cultural traditions and continues to serve as a primary healthcare

system for nearly 70–80% of the population, particularly in rural communities where access to modern healthcare facilities is limited. This widespread use of plant-based remedies highlights the urgent need for scientific validation through pharmacognosy and phytochemical sciences.

Pharmacognosy involves the identification, authentication, and quality control of crude drugs derived from natural sources such as plants, animals, and minerals. It provides essential parameters including macroscopic and microscopic characteristics, ash values, extractive values, and moisture content, which are crucial for detecting adulteration and ensuring consistency in herbal formulations. On the other hand, phytochemistry focuses on the isolation, characterization, and biological evaluation of secondary metabolites such as alkaloids, flavonoids, terpenoids, tannins, and glycosides, which are responsible for therapeutic effects.

In Nigeria, commonly used medicinal plants include *Vernonia amygdalina* (bitter leaf), widely used for treating malaria and diabetes; *Azadirachta indica* (neem), known for its antimicrobial and anti-inflammatory properties; and *Ocimum gratissimum* (scent leaf), traditionally used for respiratory and gastrointestinal disorders. Despite their widespread usage, many of these plants lack standardized pharmacognostic profiles and comprehensive phytochemical validation.

The global resurgence of interest in natural products and herbal medicines has further emphasized the importance of integrating traditional knowledge with modern scientific approaches. The World Health Organization has also encouraged the standardization and regulation of herbal medicines to ensure their safety, efficacy, and quality. Therefore, this study aims to provide a detailed pharmacognostic and phytochemical evaluation of selected Nigerian medicinal plants, contributing to their scientific validation and potential drug development.

LITERATURE REVIEW

Extensive research has been conducted on medicinal plants across Africa, with Nigeria emerging as a significant contributor due to its diverse flora. Previous studies have demonstrated that Nigerian medicinal plants contain a wide range of bioactive compounds responsible for their pharmacological activities.

Vernonia amygdalina has been extensively studied for its antidiabetic, anticancer, and antioxidant properties. The presence of sesquiterpene lactones, flavonoids, and phenolic

compounds contributes to its ability to reduce oxidative stress and regulate blood glucose levels. Several *in vivo* studies have confirmed its hypoglycemic effects, making it a promising candidate for diabetes management.

Similarly, *Azadirachta indica* has gained global recognition for its therapeutic potential. Phytochemical investigations have identified compounds such as azadirachtin, nimbin, and quercetin, which exhibit antimicrobial, antiviral, and anti-inflammatory properties. Neem extracts have been widely used in the treatment of skin infections, dental disorders, and immune-related conditions.

Ocimum gratissimum is another important medicinal plant known for its essential oil composition, particularly eugenol, thymol, and other volatile compounds. These constituents contribute to its antibacterial, antifungal, and anti-inflammatory activities. Studies have also reported its effectiveness in treating gastrointestinal disorders and respiratory infections.

Despite these advancements, the literature reveals several gaps. Many studies focus primarily on pharmacological activities without adequate pharmacognostic standardization.

Additionally, variations in phytochemical composition due to environmental factors such as soil type, climate, and harvesting time have not been sufficiently addressed. There is also a lack of comprehensive studies integrating both pharmacognostic and phytochemical analyses, which are essential for developing standardized herbal formulations.

MATERIALS AND METHODS

Plant Collection and Authentication

Fresh leaves of *Vernonia amygdalina*, *Azadirachta indica*, and *Ocimum gratissimum* were collected from different ecological regions of Nigeria, including forest and savannah zones. The plants were authenticated by a taxonomist, and voucher specimens were deposited in a recognized herbarium for future reference.




Figure 1: Map of Nigeria illustrating major ecological zones—Sahel savannah, Sudan savannah, Guinea savannah, forest zone, and mangrove swamp—along with representative regions selected for medicinal plant collection.

Sample Preparation

The collected plant materials were washed thoroughly with distilled water to remove dirt and contaminants. They were then shade-dried at room temperature for 10–14 days to preserve phytoconstituents. The dried samples were pulverized into coarse powder using a mechanical grinder and stored in airtight containers.

Pharmacognostic Evaluation

Macroscopic Analysis

The organoleptic properties such as color, odor, taste, size, and texture were examined using standard procedures.

Microscopic Analysis

Thin transverse sections of leaves were prepared using a microtome. The sections were stained with suitable reagents such as safranin and fast green to highlight cellular structures. Observations were made under a compound microscope to identify features such as stomata type, trichomes, vascular bundles, and epidermal cells.

Physicochemical Parameters

Moisture content: Determined using the loss on drying method

Total ash value: Indicates total inorganic content

Acid-insoluble ash: Measures silica content

Water and alcohol extractive values: Reflect the presence of polar and non-polar constituents

Phytochemical Screening

Qualitative phytochemical analysis was carried out using standard protocols:

Alkaloids: Detected using Mayer's and Dragendorff's reagents

Flavonoids: Identified by the Shinoda test

Tannins: Detected using ferric chloride test

Saponins: Confirmed by frothing test

Glycosides: Identified using Keller-Killiani test




Figure 2: Flowchart illustrating the overall methodology adopted for pharmacognostic evaluation and phytochemical screening of selected medicinal plants, including plant collection, sample preparation, macroscopic and microscopic analysis, extraction, qualitative phytochemical tests, and data interpretation.

RESULTS AND DISCUSSION

Pharmacognostic Findings

The macroscopic evaluation revealed distinct characteristics for each plant species. *Vernonia amygdalina* leaves were dark green with a bitter taste, *Azadirachta indica* exhibited a characteristic pungent odor, while *Ocimum gratissimum* had a strong aromatic fragrance due to essential oils.

Microscopic examination showed:

Presence of anomocytic stomata in *Vernonia amygdalina*

Multicellular trichomes in *Ocimum gratissimum*

Well-defined vascular bundles in all samples

These features are crucial for plant identification and help prevent adulteration in herbal drug preparation.

Physicochemical analysis indicated acceptable ranges of moisture content, ensuring minimal microbial growth. Ash values were within standard limits, indicating low contamination with inorganic matter.

Phytochemical Findings

Qualitative analysis confirmed the presence of multiple bioactive compounds:

Table: 1

Phytochemical	<i>Vernonia amygdalina</i>	<i>Azadirachta indica</i>	<i>Ocimum gratissimum</i>
Alkaloids	Present	Present	Present
Flavonoids	Strong	Moderate	Strong
Tannins	Moderate	Strong	Moderate
Saponins	Present	Present	Present
Glycosides	Present	Present	Present

The presence of these compounds explains the therapeutic uses of these plants:

Alkaloids contribute to antimicrobial and analgesic properties

Flavonoids act as antioxidants, protecting against oxidative stress

Tannins exhibit anti-inflammatory and antimicrobial activities

Saponins enhance immune response and possess cholesterol-lowering effects

DISCUSSION

The results strongly support the traditional use of these plants in Nigerian medicine. The combination of pharmacognostic and phytochemical analyses provides a reliable approach for standardizing herbal drugs. However, variations in phytochemical composition were observed, which may be attributed to environmental factors such as soil composition, rainfall, and harvesting time.

These findings emphasize the need for:

- Standardized cultivation practices
- Controlled harvesting conditions
- Advanced analytical techniques such as HPLC and GC-MS for compound identification

The integration of traditional knowledge with scientific validation not only enhances drug discovery but also ensures the safety and efficacy of herbal medicines on a global scale.

CHALLENGES IN PHARMACOGNOSY AND PHYTOCHEMICAL RESEARCH

Pharmacognosy and phytochemical sciences in Nigeria hold immense promise for drug discovery and validation of traditional medicine; however, several critical challenges continue to hinder their full development and global acceptance. These challenges span scientific, infrastructural, regulatory, and socio-cultural dimensions.

1. Lack of Standardization of Herbal Drugs

One of the most significant challenges is the absence of uniform standards for the identification, processing, and formulation of herbal medicines. Unlike synthetic drugs, which undergo strict quality control, herbal products often vary in composition due to differences in plant species,

geographical origin, harvesting time, and processing methods. This variability leads to inconsistent therapeutic outcomes and raises concerns about safety and efficacy.

Pharmacognostic parameters such as ash values, extractive values, and microscopic characteristics are not consistently applied across studies or commercial preparations. The lack of standardized protocols makes it difficult to establish reproducibility and reliability in research findings.

2. Variability in Phytochemical Composition

The phytochemical composition of medicinal plants is highly influenced by environmental factors such as soil type, climate, altitude, rainfall, and seasonal variations. For example, the concentration of alkaloids or flavonoids in a plant harvested during the rainy season may differ significantly from those collected in the dry season.

This variability creates challenges in:

- Ensuring consistent bioactive compound concentration
- Reproducing experimental results
- Developing standardized herbal formulations

Without controlled cultivation and harvesting practices, it becomes difficult to guarantee the quality of plant-derived drugs.

3. Limited Access to Advanced Analytical Technologies

Advanced analytical techniques such as High-Performance Liquid Chromatography (HPLC), Gas Chromatography-Mass Spectrometry (GC-MS), and Nuclear Magnetic Resonance (NMR) spectroscopy are essential for identifying and characterizing bioactive compounds. However, many research institutions in Nigeria face limitations due to:

- High cost of equipment procurement and maintenance
- Inadequate funding for scientific research
- Lack of trained personnel to operate sophisticated instruments

As a result, many studies rely only on preliminary qualitative tests rather than detailed compound identification, limiting the depth and global competitiveness of research outputs.

4. Inadequate Documentation of Traditional Knowledge

Traditional medicinal knowledge in Nigeria is often transmitted orally from one generation to another. This lack of written documentation poses a risk of knowledge loss, especially as older practitioners pass away without transferring their expertise.

5. Regulatory and Quality Control Challenges

The regulation of herbal medicines in Nigeria is still evolving. Although agencies like the National Agency for Food and Drug Administration and Control (NAFDAC) oversee drug approval, enforcement remains inconsistent.

Key issues include:

- Presence of adulterated or contaminated herbal products in the market
- Lack of standardized guidelines for herbal drug approval
- Limited post-marketing surveillance

These regulatory gaps reduce public confidence and hinder international acceptance of Nigerian herbal products.

6. Contamination and Adulteration Risks

Medicinal plants are susceptible to contamination from heavy metals, pesticides, microbial toxins, and environmental pollutants. In some cases, intentional adulteration occurs when cheaper or incorrect plant species are substituted to increase profit margins.

This leads to:

- Reduced therapeutic efficacy
- Potential toxicity and adverse health effects
- Difficulty in authentication of plant materials

Pharmacognostic techniques such as microscopy and physicochemical analysis are essential but are not always rigorously applied.

7. Limited Clinical Validation

While many Nigerian medicinal plants have demonstrated promising results in laboratory

studies, there is a lack of well-designed clinical trials to confirm their safety and efficacy in humans. Most research remains at the in vitro or animal study level.

Challenges in clinical validation include:

- High cost of clinical trials
- Ethical and regulatory complexities
- Limited collaboration between academia and healthcare institutions

Without clinical evidence, it is difficult for herbal medicines to gain acceptance in mainstream healthcare systems.

8. Insufficient Funding and Research Support

Research in pharmacognosy and phytochemistry requires substantial financial investment for fieldwork, laboratory analysis, and publication. However, funding in Nigeria is often limited, leading to:

- Small-scale studies with limited impact
- Inability to acquire modern equipment
- Reduced participation in international research collaborations

This financial constraint significantly slows down scientific progress in the field.

9. Lack of Interdisciplinary Collaboration

Pharmacognosy and phytochemical research require collaboration among botanists, chemists, pharmacologists, and medical professionals. In many cases, such interdisciplinary collaboration is weak or absent, resulting in fragmented research efforts.

An integrated approach is necessary to:

- Fully understand plant bioactivity
- Develop effective herbal formulations
- Translate research findings into clinical applications

10. Global Competition and Acceptance Barriers

Despite the rich medicinal plant resources in Nigeria, herbal products face challenges in

entering the global market due to:

- compliance with international quality standards (e.g., WHO, FDA guidelines)
- Lack of standardized documentation and clinical data
- Competition from well-established herbal industries in countries like China and India

To overcome these barriers, Nigerian research must align with global best practices in pharmacognosy and phytochemical sciences.

CONCLUSION

The present study provides a comprehensive pharmacognostic and phytochemical evaluation of selected Nigerian medicinal plants, emphasizing their scientific relevance in modern drug discovery and development. The detailed macroscopic, microscopic, and physicochemical assessments established reliable identification parameters that are essential for ensuring the authenticity and quality of crude drugs. These pharmacognostic standards serve as a critical foundation for preventing adulteration and maintaining consistency in herbal formulations.

The phytochemical screening revealed the presence of key bioactive constituents such as alkaloids, flavonoids, tannins, saponins, and glycosides across the studied plant species. These compounds are widely recognized for their therapeutic properties, including antimicrobial, antioxidant, anti-inflammatory, and immunomodulatory activities. The findings strongly validate the traditional use of these plants in Nigerian ethnomedicine and highlight their potential as sources of novel pharmacologically active agents.

However, the study also underscores several challenges, including variability in phytochemical composition, lack of standardization, limited access to advanced analytical tools, and insufficient clinical validation. Addressing these limitations is essential for enhancing the credibility, safety, and global acceptance of herbal medicines. Strengthening regulatory frameworks, improving research funding, and fostering interdisciplinary collaboration will play a crucial role in advancing pharmacognosy and phytochemical sciences in Nigeria.

In a broader context, the integration of traditional medicinal knowledge with modern scientific approaches offers a promising pathway for sustainable healthcare solutions. With continued

research and standardization, Nigerian medicinal plants have the potential to contribute significantly to global pharmaceutical development and natural product-based therapeutics.

REFERENCES

1. Sofowora, A. (2008). *Medicinal Plants and Traditional Medicine in Africa* (3rd ed.). Spectrum Books Limited.
2. Harborne, J. B. (1998). *Phytochemical Methods: A Guide to Modern Techniques of Plant Analysis* (3rd ed.). Chapman & Hall.
3. Edeoga, H. O., Okwu, D. E., & Mbaebie, B. O. (2005). Phytochemical constituents of some Nigerian medicinal plants. *African Journal of Biotechnology*, 4(7), 685–688.
4. Odugbemi, T. (2006). *Outlines and Pictures of Medicinal Plants from Nigeria*. University of Lagos Press.
5. Trease, G. E., & Evans, W. C. (2009). *Pharmacognosy* (16th ed.). Saunders Elsevier.
6. Kokate, C. K., Purohit, A. P., & Gokhale, S. B. (2010). *Pharmacognosy* (45th ed.). Nirali Prakashan.
7. World Health Organization. (2013). WHO traditional medicine strategy 2014–2023. <https://www.who.int>
8. Ajayi, I. A., Ajibade, O., & Oderinde, R. A. (2011). Preliminary phytochemical analysis of some plant seeds. *Research Journal of Chemical Sciences*, 1(3), 58–62.
9. Okwu, D. E. (2004). Phytochemicals and vitamin content of indigenous spices of Southeastern Nigeria. *Journal of Sustainable Agriculture and Environment*, 6(1), 30–37.
10. Ncube, N. S., Afolayan, A. J., & Okoh, A. I. (2008). Assessment techniques of antimicrobial properties of natural compounds. *African Journal of Biotechnology*, 7(12), 1797–1806.
11. Cowan, M. M. (1999). Plant products as antimicrobial agents. *Clinical Microbiology Reviews*, 12(4), 564–582.
12. Fabricant, D. S., & Farnsworth, N. R. (2001). The value of plants used in traditional medicine. *Environmental Health Perspectives*, 109(Suppl 1), 69–75.
13. Gurib-Fakim, A. (2006). Medicinal plants: Traditions of yesterday and drugs of tomorrow. *Molecular Aspects of Medicine*, 27(1), 1–93.

Cite as:

Dr. Chinedu Okafor, Amina Bello, Ibrahim Musa (2026). Phytochemical Investigation and Pharmacognostic Evaluation of Selected Medicinal Plants in Nigeria. International Journal of Pharmacognosy and Phytochemical Sciences, 8(1), 20-32.

<https://doi.org/10.5281/zenodo.19707600>