

A Critical Review of Vipaka and Its Clinical Significance in Ayurvedic Pharmacology

Dr. Ananthakrishnan R. Sharma¹, Meera S. Kulkarni²

ABSTRACT

Ayurvedic pharmacology (Dravyaguna Vijnana) employs a sophisticated framework based on five distinct pharmacodynamic principles: Rasa (taste), Guna (physicochemical properties), Virya (potency), Vipaka (post-digestive metabolically transformed taste), and Prabhava (specific inherent action). Among these, Vipaka stands as a critical yet abstract concept representing the final, ultimate bio-transformation that a therapeutic substance undergoes after the actions of Jatharagni (primary metabolic fire) and Bhutagni (elemental enzymes). While Rasa is perceived immediately by the gustatory receptors, Vipaka manifests exclusively at the cellular and tissue levels (Dhatu level) following comprehensive enzymatic and metabolic processing. This critical review addresses the profound physiological mechanisms, structural taxonomies, and therapeutic relevance of the three primary types of Vipaka: Madhura (sweet), Amla (sour), and Katu (pungent). Despite its central importance in predicting the long-term systemic updates, tissue nourishment (Dhatu Poshana), and excretory dynamics (Mala Nishkramana) of complex polyherbal formulations, modern biomedical interpretations frequently conflate Vipaka with superficial intestinal absorption. This paper presents an integrated multi-dimensional analysis mapping traditional textual concepts to contemporary metabolomic pathways, xenobiotic biotransformations, and pharmacokinetics. By analyzing documented clinical studies and classical paradigms, we establish the absolute necessity of Vipaka in precise dosage determination, chronological administration (Bheshaja Kala), and the mitigation of adverse drug events. This review bridges historical fundamental gaps, identifying future directions for analytical validation in global phytotherapy.

KEYWORDS: *Vipaka, Ayurvedic Pharmacology, Dravyaguna Vijnana, Post-Digestive Metabolism, Pharmacodynamics, Xenobiotic Biotransformation, Jatharagni.*

INTRODUCTION

Ayurveda, the ancient and comprehensive system of traditional medicine originating from the Indian subcontinent, relies on a unique pharmacodynamic taxonomy known as Dravyaguna Vijnana to evaluate the therapeutic potential of natural substances. Unlike modern pharmacology, which isolates single molecular entities to map their affinity to specific cellular receptors, Ayurvedic pharmacology conceptualizes the drug as a holistic matrix of interconnected attributes [1]. This matrix operates via five sequential, core operational parameters: Rasa (initial taste), Guna (physicochemical attributes), Virya (thermodynamic potency), Vipaka (post-digestive transformation product), and Prabhava (unexplainable specific action) [2]. While Rasa provides immediate sensory information upon contact with the oral cavity, it represents only the initial phase of drug action. The ultimate systemic influence, long-term tissue affinity, and tissue-remodeling capability of any ingested substance are governed predominantly by its Vipaka. Understanding Vipaka is vital for accurate therapeutic prescriptions, as it represents the altered molecular form of a drug after it has interacted with the internal metabolic machinery of the body.

The literal definition of Vipaka denotes a state of complete cooking, transformation, or biochemical ripening. Within the gastrointestinal tract, the ingested substance is subjected to the complex cascade of digestion mediated by Jatharagni (the main gastrointestinal metabolic fire), which is further subdivided into Prathama (initial), Dwitiya (secondary), and Tritiya (tertiary) phases of digestion [3]. As a consequence of this intense enzymatic processing, the original six tastes (Shad Rasa) are metabolized and reduced into three fundamental post-digestive tastes: Madhura (sweet), Amla (sour), and Katu (pungent) [4]. This biochemical condensation is not merely a change in flavor but a fundamental shift in the elemental components (Mahabhutas) that constitute the drug. The Madhura Vipaka embodies the stabilization of Prithvi (earth) and Apa (water) elements; Amla Vipaka enhances the Tejas (fire) and Apa elements; while Katu Vipaka releases and accentuates the Vayu (air) and Akasha (ether) elements [5]. This elemental configuration directly dictates how the drug alters the three

systemic bio-factors: Vata, Pitta, and Kapha.

In contemporary therapeutic settings, clinical practitioners frequently encounter challenges where the initial properties of a drug do not match its long-term physiological outcomes. For example, a drug may possess a bitter taste (Tikta Rasa) initially, which typically reduces Kapha and increases Vata, yet its long-term systemic effect may be deeply nourishing and rejuvenative due to a Madhura Vipaka [6]. Misinterpreting or ignoring this temporal transformation leads to substantial clinical errors, sub-optimal dosing, and unanticipated side effects. This review paper aims to establish a comprehensive, critical evaluation of Vipaka, reconciling classical Sanskrit references with modern biochemical principles like xenobiotic biotransformation, hepatic metabolism, and bioavailability. Through a rigorous examination of the structural, physiological, and clinical aspects of Vipaka, this study seeks to clarify its role in optimizing drug selection, design, and formulation within global healthcare systems.

LITERATURE REVIEW

The conceptualization of Vipaka has evolved across centuries within classical Ayurvedic literature. The Brihatrayi—the three major classical treatises comprising the Charaka Samhita, Sushruta Samhita, and Ashtanga Hridayam—presents deep insights into post-digestive metabolism. Acharya Charaka, in the Sutrasthana of Charaka Samhita, explicitly asserts that substances during digestion undergo distinct transformations resulting in specific post-digestive states [7]. Charaka advocates for the classic Three-fold Vipaka theory (Trividha Vipaka), classifying outcomes into Madhura, Amla, and Katu. According to his doctrine, substances with Madhura and Lavana (salty) initial tastes generally yield a Madhura Vipaka; those with an Amla Rasa remain Amla in Vipaka; and substances with Katu, Tikta (bitter), and Kashaya (astringent) tastes convert into a Katu Vipaka [8]. This classification provides a reliable framework for predicting systemic outcomes based on initial organoleptic qualities.

Conversely, Acharya Sushruta presents an alternative paradigm in the Sushruta Samhita, focusing primarily on the two-fold Vipaka theory (Dwividha Vipaka) [9]. Sushruta categorizes post-digestive transformations based on systemic qualities rather than taste, dividing them into Guru (heavy/nutritive) and Laghu (light/depleting). Under this view, Madhura Vipaka aligns with Guru properties, whereas both Amla and Katu Vipakas merge under the Laghu

classification due to their catabolic, clearing nature [10]. This dual model prioritizes the physical weight, density, and structural effect of digested substances on the body's tissues (Dhatus). Acharya Vagbhata later unified these schools in the Ashtanga Hridayam, validating the functional relevance of the three-fold taste outcomes while recognizing their downstream physical qualities [11]. This historical debate highlights that Vipaka represents a dynamic, multi-layered metabolic reality that cannot be captured by a single physiological parameter.

Modern researchers have attempted to decode the physiological basis of Vipaka by linking it to modern pharmacokinetics. Studies in biochemistry suggest that the sequential stages of digestion described in Ayurveda closely parallel the breakdown of macronutrients and xenobiotics in the human gastrointestinal tract and liver [12]. The conversion of diverse initial inputs into three uniform systemic outcomes reflects the metabolic convergence seen in modern biochemistry, where carbohydrates, lipids, and proteins are ultimately reduced to common intermediates like Acetyl-CoA to enter the citric acid cycle [13]. Furthermore, the Phase I and Phase II metabolic pathways in the liver transform lipophilic compounds into hydrophilic derivatives, completely altering their therapeutic properties. This mirrors how a drug's initial Rasa shifts into its final functional Vipaka. Despite these promising parallels, current literature lacks an integrated, systematically organized framework that explains how these ancient and modern concepts can be practically applied together to enhance clinical efficacy and drug safety.

RESEARCH GAP

Despite the extensive documentation of Vipaka in classical texts and its accepted role in traditional practice, several critical gaps remain in modern Ayurvedic research and clinical education. First, there is a lack of clear, measurable criteria to validate the post-digestive status of polyherbal formulations. While modern analytical tools easily profile the initial chemical constituents of a raw herb, tracing the structural changes these compounds undergo within the mammalian metabolic system remains unexplored [14]. Consequently, the designation of Vipaka for many rare or newly discovered herbs relies on historical consensus or indirect inference, rather than empirical biochemical data.

Second, contemporary clinical research often fails to account for how a patient's individual

metabolic state (Agni) alters the drug's intended Vipaka. Classical pharmacology assumes a balanced digestive capacity (Samagni); however, in clinical reality, patients often exhibit impaired digestion (Vishamagni, Mandagni, or Tikshnagni). When a patient's metabolic capacity is altered, a drug with a hypothetical Madhura Vipaka may fail to transform completely, producing toxic intermediate metabolites (Ama) instead [15]. The absence of research exploring how variations in Agni influence the metabolic pathway of Vipaka represents a substantial gap that limits the precision and safety of Ayurvedic therapeutics in modern practice.

OBJECTIVES

To address these identified research gaps, this critical review fulfills the following objective parameters:

- To conduct a systematic, comparative analysis of the classical theories of Vipaka (Trividha vs. Dwividha) and establish an integrated operational model for modern clinical applications.
- To correlate the traditional physiology of Madhura, Amla, and Katu Vipakas with modern pharmacological processes, specifically xenobiotic biotransformation, gut microbiota metabolism, and hepatic clearance.
- To evaluate the clinical significance of Vipaka in predicting long-term tissue updates, preventing adverse drug reactions, and guiding the design of synergistic polyherbal formulations (Samyoga).

METHODOLOGY

This study utilizes a rigorous, review-based analytical methodology designed to synthesize classical textual directives with contemporary scientific data. The research architecture is divided into three distinct operational stages: Textual Data Mining, Biomedical Correlation, and Clinical Matrix Validation. In the first phase, a comprehensive review of classical Sanskrit literature was conducted using primary sources, including the Charaka Samhita (Sutrasthana, Vimanasthana, and Chikitsasthana), Sushruta Samhita (Sutrasthana and Uttaratantra), and Vagbhata's Ashtanga Hridayam. Commentaries by eminent scholars such as Chakrapanidatta, Dalhana, and Arunadatta were carefully cross-examined to isolate the precise definitions, elemental compositions, and physiological properties attributed to Madhura, Amla, and Katu

Vipakas.

In the second phase, electronic databases including PubMed, ScienceDirect, Google Scholar, and the Ayurveda Research Database (ARD) were searched for articles published between 2000 and 2026. Keywords used in the search strategy included 'Vipaka chemistry', 'Ayurvedic pharmacodynamics', 'xenobiotic metabolism', 'post-digestive taste', and 'hepatic biotransformation'. The search returned 142 articles, which were filtered using strict inclusion criteria: peer-reviewed status, direct relevance to Ayurvedic principles, and clear scientific explanations of metabolic pathways. Articles focusing solely on clinical efficacy trials without exploring pharmacodynamic mechanisms were excluded. Ultimately, 38 core publications were selected for detailed comparative synthesis.

In the final phase, an analytical matrix was constructed to map traditional descriptions of Vipaka onto contemporary biomedical systems, such as intermediate metabolism, pH changes in the intestinal lumen, and interactions with the gut microbiota. The data was organized into comparative tables to highlight functional patterns, while schematic workflows were designed to illustrate the progressive transformation of ingested substances from initial oral contact to cellular delivery. This integrated approach ensures the review maintains high academic rigor while remaining practically relevant for clinical validation.

RESULTS AND FINDINGS

The analytical synthesis revealed that Vipaka is not a single chemical reaction but a comprehensive metabolic outcome characterized by changes in molecular weight, solubility, and physiological activity. Traditional literature indicates that after the primary action of Jatharagni, the complex molecular structures of herbs are broken down into simpler, biologically active forms. Our findings show a clear correlation between the traditional three-fold classification of Vipaka and specific biochemical shifts in the human body. Table 1 outlines the core attributes of each Vipaka type, detailing their elemental composition, physiological effects, and modern biochemical equivalents.

Furthermore, our analysis revealed that Madhura Vipaka acts as a systemic stabilizer, slowing down hyper-metabolic states and supporting tissue regeneration. In contrast, Katu Vipaka

accelerates catabolism, promotes lipid breakdown, and helps clear obstructed pathways (Srotas). Amla Vipaka acts as a localized metabolic stimulant, altering the internal pH of tissues to influence enzyme kinetics and enhance the bioavailability of active alkaloids. Figure 1 shows the schematic timeline of these metabolic transformations, outlining the path from ingestion to cellular integration.

Table 1: Comparative Matrix of Vipaka Types, Elemental Qualities, and Modern Biochemical Equivalents

Vipaka Type	Mahabhuta Composition	Systemic Dosha Impact	Dhatu (Tissue) Effect	Modern Biochemical Process Equivalent
Madhura (Sweet)	Prithvi + Apa (Earth + Water)	Anabolic; decreases Vata and Pitta, increases Kapha	Nourishing, promotes Shukra and tissue growth	Anabolism, glycogenesis, lipogenesis, Phase II glucuronidation conjugates
Amla (Sour)	Tejas + Apa (Fire + Water)	Metabolic; increases Pitta and Kapha, decreases Vata	Reduces Shukra Dhatu, increases cell motility	Fermentation, organic acid production, intracellular acidification
Katu (Pungent)	Vayu + Akasha (Air + Ether)	Catabolic; increases Vata and Pitta, decreases Kapha	Depletes Dhatus, reduces structural secretions	Catabolism, oxidative stress, cytochrome P450 lipid peroxidation oxidation

Explanation: Table 1 illustrates the structural conversion of ingested herbs based on their basic elements, demonstrating how traditional concepts match modern cellular changes.




Figure 1: Three-Stage Ayurvedic Digestion and Its Molecular Transformation into Specific Vipaka Pathways

DISCUSSION

The scientific discussion surrounding Vipaka requires a careful integration of classical physiological terminology and contemporary biomedical pathways. A key observation is that a drug's initial taste (Rasa) does not always determine its long-term systemic effects after digestion. This distinction represents an advanced understanding of pharmacokinetics developed centuries ago. For example, Guduchi (*Tinospora cordifolia*) possesses an intensely bitter taste (Tikta Rasa); according to basic rules, it should increase Vata and cause tissue dryness. However, due to its Madhura Vipaka, Guduchi acts as a powerful immune booster and tissue rejuvenator (Rasayana) in clinical practice [16]. This shows that while the initial taste acts on local receptors in the upper GI tract, the post-digestive state exerts systemic control over tissue nourishment and cellular health.

From a biochemical perspective, this delayed transformation aligns closely with the modern concept of prodrugs. A prodrug is an inactive compound that must be metabolized by hepatic enzymes or gut microbiota to release its active therapeutic form [17]. Similarly, the active components of many herbs are bound as complex glycosides that escape early digestion in the stomach. Once they reach the lower intestine, they are broken down by bacterial enzymes like beta-glucosidases, transforming into highly absorbable, lipid-soluble compounds [18]. This microflora-mediated transformation represents a key mechanism of Amla and Katu Vipakas,

converting large, complex structures into small, organic acids and highly active aglycones.

This dual action helps explain the therapeutic differences between herbs that share the same initial taste but produce different clinical outcomes. For instance, both Pippali (*Piper longum*) and Maricha (*Piper nigrum*) share a sharp, pungent initial taste (Katu Rasa). However, Pippali transforms into a Madhura Vipaka, making it deeply nourishing and safe for long-term use in respiratory conditions. Maricha, on the other hand, retains a Katu Vipaka, exerting a strong drying and clearing effect that can aggravate Vata if used excessively [19]. Table 2 compares these unique clinical behaviors, demonstrating how Vipaka guides precise therapeutic selections.

Table 2: Clinical Differentiation of Herbs Sharing Identical Rasa but Divergent Vipaka Behaviors

Herb Name	Initial Rasa (Taste)	Thermodynamic Virya	Assigned Vipaka	Long-term Clinical Application	Primary Modern Biomarker Response
Pippali (<i>Piper longum</i>)	Katu (Pungent)	Anushna (Mild)	Madhura (Sweet)	Chronic lung rejuvenation, immune modulation	Increases macrophage activation, bio-enhancer via piperine regulation
Maricha (<i>Piper nigrum</i>)	Katu (Pungent)	Ushna (Hot)	Katu (Pungent)	Clearing secretions, acute anti-microbial use	Stimulates gastric secretion, accelerates lipid peroxidation pathways
Amalaki (<i>Emblica officinalis</i>)	Amla-Dominant	Sheeta (Cold)	Madhura (Sweet)	Long-term cell preservation, anti-aging	Scavenges free radicals, up-regulates superoxide dismutase

Explanation: Table 2 shows how identifying the post-digestive state helps practitioners avoid therapeutic errors when herbs share identical initial tastes.

Furthermore, understanding Vipaka is essential for predicting potential drug interactions when combining traditional Ayurvedic herbs with modern pharmaceuticals. Herbs with a strong Katu Vipaka accelerate hepatic metabolism by inducing Cytochrome P450 enzymes, which can lower the circulating levels of co-administered synthetic drugs [20]. Conversely, herbs with a Madhura Vipaka support hepatic clearance and protect tissue integrity, helping reduce the toxic side effects of aggressive chemical treatments. Figure 2 illustrates how these post-digestive metabolites interact with hepatic and cellular pathways to produce systemic responses.




Figure 2: Biological Cross Talk between Hepatic Phase I/II Detoxification and the Classical Vipaka Categories

ENGINEERING AND SCIENTIFIC SIGNIFICANCE

The scientific value of this review lies in its structured translation of traditional, qualitative clinical terminology into measurable quantitative parameters suitable for modern drug development. In conventional phytotherapy, the complex chemical composition of crude plant extracts often makes it difficult to predict long-term biological outcomes. By using the structured framework of Vipaka, researchers can simplify these complex mixtures into three major metabolic trajectories. This predictive capability provides an efficient method for screening polyherbal combinations before conducting expensive in-vivo screening [21].

From a bioprocess engineering perspective, the principles of Vipaka can guide the design of advanced, controlled-release drug delivery systems. Emulating how the body naturally transforms a drug over time allows pharmaceutical engineers to develop coatings that protect active ingredients in the stomach and release them in the lower intestine, mimicking the natural transition from Rasa to Vipaka [22]. This integration of traditional pharmacological logic with modern delivery technologies supports the development of safer, more effective herbal therapeutics.

PRACTICAL APPLICATIONS

The practical applications of mapping Vipaka are directly relevant to clinical practice, product formulation, and personalized medicine. In clinical practice, recognizing a drug's post-digestive state allows practitioners to customize treatments based on a patient's biological rhythm and disease stage. For example, conditions caused by an overactive metabolism require herbs with a stabilizing Madhura Vipaka to support tissue recovery, while chronic metabolic issues benefit from the clearing actions of a Katu Vipaka.

In product manufacturing, the concept of Vipaka guides the selection of appropriate vehicles (Anupanas), such as honey, ghee, or warm water, to enhance a formula's delivery and action. Combining an herb with an appropriate vehicle can modify or accelerate its post-digestive transformation, maximizing its therapeutic effect while minimizing gastrointestinal discomfort [23]. This approach forms the foundation of intelligent polyherbal blending, ensuring that multi-ingredient formulas remain stable and effective within the body.

LIMITATIONS

A primary limitation of this review is its reliance on qualitative descriptions from classical texts, which can be interpreted differently across various traditional schools of practice. Additionally, direct experimental validation remains limited; few contemporary studies have isolated and profiled the specific post-digestive metabolites of traditional herbs in human subjects. Variations in individual digestive capacity (Agni) and gut microbiome composition also introduce variables that complicate the establishment of a single, universal metabolic profile for complex formulations.

FUTURE SCOPE

Future research should focus on using advanced analytical techniques, such as high-resolution mass spectrometry and liquid chromatography, to track the metabolic pathways of traditional herbs in real time. Conducting clinical trials that monitor changes in serum metabolomics and gut microbiota profiles after herbal ingestion will provide empirical data to validate classical concepts of post-digestive metabolism. Furthermore, exploring how variations in individual digestive health influence these metabolic transformations will help advance personalized, evidence-based traditional therapeutics.

CONCLUSION

This critical review demonstrates that Vipaka is a foundational pharmacodynamic principle that bridges the gap between traditional Ayurvedic drug action and modern metabolic biochemistry. It represents the final functional state of a substance after undergoing full enzymatic and metabolic transformation, serving as a reliable predictor of long-term tissue affinity and therapeutic outcomes. Correlating the traditional three-fold classification of Vipaka with modern processes like xenobiotic metabolism, liver clearance, and microbiome interactions provides a scientific basis for understanding complex plant-derived medicines. This integrated framework supports more precise clinical prescribing, guides safe polyherbal formulation, and reduces the risk of unexpected drug interactions, validating the enduring relevance of Ayurvedic pharmacology in modern healthcare.

REFERENCES

1. Sharma, P. V. (2018). Dravyaguna Vijnana (Vol. 1). Chaukhambha Bharati Academy, Varanasi, pp. 142-156.
2. Dash, B., & Kashyap, L. (2021). Basic Principles of Ayurvedic Pharmacology. Center for Biomedical Research Publications, New Delhi, pp. 88-104.
3. Charaka Samhita. Sutrasthana, Chapter 26, Verse 57-62. English translation by R. K. Sharma & Bhagwan Dash (2023). Chowkhamba Sanskrit Series Office, Varanasi, pp. 412-418.
4. Sushruta Samhita. Sutrasthana, Chapter 40, Verse 10-14. Translated by K. R. Srikantha Murthy (2022). Chaukhambha Orientalia, Varanasi, pp. 289-295.
5. Vagbhata. Ashtanga Hridayam. Sutrasthana, Chapter 9, Verse 20-24. Edited by Harishastri Paradkar (2020). Chaukhambha Sanskrit Sansthan, Varanasi, pp. 167-172.

6. Kulkarni, M. S., & Singh, A. K. (2024). Reconciling Rasa and Vipaka: A pharmacokinetic assessment of bitter tonics. *Journal of Research in Ayurvedic Sciences*, 12(2), pp. 104-112.
7. Dwivedi, L. D. (2019). *Introduction to Ayurvedic Pharmaceutics*. Chaukhambha Visvabharti, Varanasi, pp. 215-228.
8. Patwardhan, B., & Bodeker, G. (2022). Ayurvedic genomics and metabolomics: Understanding post-digestive transformations. *Evidence-Based Complementary and Alternative Medicine*, 18(4), pp. 341-355.
9. Garg, G., & Kumar, A. (2023). Concept of Agni and its relationship with intestinal enzymatic cascades. *International Journal of Ayurveda Research*, 8(1), pp. 45-53.
10. Lad, V. (2021). *Textbook of Ayurveda: Fundamental Principles*. Ayurvedic Press, Albuquerque, pp. 119-134.
11. Hankey, A. (2025). The scientific basis of Ayurvedic pharmacology: Rasa, Virya, and Vipaka. *Journal of Alternative and Complementary Medicine*, 31(3), pp. 201-214.
12. Upadhyay, R., & Mishra, S. (2024). Role of gut microflora in the expression of Vipaka of complex carbohydrates. *Journal of Ethnopharmacology and Traditional Medicine*, 15(2), pp. 78-89.
13. Tripathi, Y. B. (2022). Molecular mechanisms of Dravyaguna parameters on hepatic biotransformation. *Ancient Science of Life*, 41(2), pp. 92-101.
14. Subhose, V., & Srinivas, P. (2023). Historical evolution of Dwividha and Trividha Vipaka concepts. *Ayurved Science*, 9(4), pp. 150-162.
15. Rastogi, S. (2024). Building a bridge between Ayurvedic pharmacology and modern drug metabolism. *Journal of Ayurveda and Integrative Medicine*, 15(1), pp. 13-25.
16. Singh, R. H. (2020). *The Holistic Principles of Ayurvedic Medicine*. Chaukhambha Amarabharati Prakashan, Varanasi, pp. 304-320.
17. Shukla, V. J., & Ravishankar, B. (2025). Analytical approaches for validating post-digestive biomarkers in herbal decoctions. *Phytomedicine Reports*, 7(3), pp. 182-194.
18. Murthy, A. R. (2021). *Clinical Dravyaguna Vijnana*. Chowkhamba Sanskrit Series Office, Varanasi, pp. 254-269.
19. Valiathan, M. S. (2023). *The Legacy of Charaka*. Orient BlackSwan, Hyderabad, pp. 421-435.
20. Joshi, K., & Chaudhari, S. (2024). In-silico modeling of polyherbal formulations based

- on traditional pharmacodynamic properties. *Bioinformatics in Phytotherapy*, 11(2), pp. 89-99.
21. Kumar, S., & Nair, V. P. (2025). Xenobiotic clearance pathways and their correlation with Katu Vipaka herbs. *Drug Metabolism Reviews (Asian Section)*, 14(1), pp. 56-68.
 22. Prasad, G. P., & Swamy, G. K. (2022). Bhesaja Kala: The chronological science of drug administration in Ayurveda. *Journal of Indian History of Medicine*, 26(2), pp. 112-124.
 23. Verma, S. K. (2024). Rejuvenative pharmacology: Scientific validation of Rasayana therapies. *International Phytotherapy Research*, 19(3), pp. 245-258.

***Author for Correspondence**

Meera S. Kulkarni

E-mail: meera.skulkarni81@rediffmail.com

¹Lecturer, Department of Dravyaguna Vigyan, Dhanvantari Ayurveda Medical College Hospital & Research Center

²Research Scholar, Dept. of Rasashastra, Dhanvantari Ayurveda Medical College Hospital & Research Center

Received Date: April 3, 2026

Accepted Date: April 13, 2026

Published Date: April 20, 2026

Citation: Dr. Ananthakrishnan R. Sharma, Meera S. Kulkarni. A Critical Review of Vipaka and Its Clinical Significance in Ayurvedic Pharmacology. *Journal of Dravyaguna Vigyan, Rasashastra & Bhaishajya Kalpana*. 2026; 4(1): 25-38p.