MANTECH Publications

Journal of Engineering Mathematics & Statistics
Volume 3, Issue 2, July-December 2019

Approximation Theory & Meshless Methods

Umesh Patil[1], Depesh Deshpande[2], Ankur. Choudhury[3]

Associate Professor[1], Students[2, 3]

Department of Applied Mathematics

North Eastern Rural University, Assam, India

Email ID: *Umeshpatil11@gmail.com[1], depeshdespandee@rediffmail.com[2], ankurchaudhary909@yahoo.com[3]*

Abstract

Approximation theory forms the mathematical backbone of many numerical and computational methods used to solve real-world problems where exact solutions are difficult or impossible to obtain. In recent decades, meshless (or meshfree) methods have emerged as a powerful alternative to classical mesh-based numerical techniques such as finite difference, finite element, and finite volume methods. These approaches rely heavily on approximation theory, particularly interpolation, regression, and functional approximation using scattered data. This paper presents a comprehensive review of approximation theory and its close connection with meshless methods. Fundamental concepts such as polynomial approximation, radial basis functions, moving least squares, and kernel-based approximations are discussed in detail. Various meshless methods, including the Element-Free Galerkin method, Smoothed Particle Hydrodynamics, and Local Petrov–Galerkin methods, are reviewed with emphasis on their theoretical foundations, advantages, and limitations. Applications in engineering, fluid dynamics, solid mechanics, and computational physics are also highlighted. The paper aims to provide a unified perspective that connects classical approximation theory with modern meshless numerical techniques, while also identifying current challenges and future research directions.

Keywords: *Approximation theory, meshless methods, radial basis functions, moving least squares, numerical analysis, scattered data approximation*

Page 104-114 © MANTECH PUBLICATIONS 2019. All Rights Reserved

INTRODUCTION

Approximation theory is a classical branch of mathematics that deals with how functions can be approximated by simpler or more tractable functions, such as polynomials, splines, or trigonometric series. The need for approximation arises naturally in science and engineering, where exact analytical solutions are often unavailable due to complex geometries, nonlinearities, or irregular boundary conditions. Over the years, approximation theory has evolved from pure mathematical interest into a practical tool that supports numerical computation and simulation.

Traditional numerical methods for solving partial differential equations (PDEs), such as the finite element method (FEM) and finite difference method (FDM), rely on predefined meshes or grids. While these methods are well established and widely used, they face certain difficulties when dealing with large deformations, moving boundaries, crack propagation, or higher-dimensional problems. Mesh generation itself can be time-consuming and error-prone, especially for complex geometries.

Meshless methods were introduced to overcome some of these limitations. Instead of relying on a mesh, these methods use a set of scattered nodes distributed over the problem domain. Approximation theory plays a central role in meshless methods, as the construction of shape functions, interpolation schemes, and numerical integration techniques all depend on approximation principles.

This paper reviews the theoretical foundations of approximation theory and explains how they are utilized in meshless methods. The discussion is kept broad enough to be accessible to researchers from applied mathematics and engineering backgrounds, while still providing sufficient depth for academic study.

Fundamentals of Approximation Theory

Approximation theory is concerned with the representation of complex mathematical objects using simpler and more manageable functions. The main objective is not to find an exact representation, but rather to achieve an approximation that is sufficiently accurate for practical computation while

Page 104-114 © MANTECH PUBLICATIONS 2019. All Rights Reserved

remaining stable and efficient. This field forms the mathematical basis for most numerical methods used in scientific computing, including meshless techniques.

Concept of Approximation

In its most basic form, approximation theory aims to replace a complicated or unknown function f(x)f(x)f(x) with a simpler function g(x)g(x)g(x), such that the difference between the two functions is minimized according to a chosen criterion. The function g(x)g(x)g(x) is usually selected from a finite-dimensional space, such as polynomials, splines, or basis functions, which makes computation feasible.

The approximation process depends strongly on how the error between f(x)f(x)f(x) and g(x)g(x)g(x) is measured. This error is defined using norms, and the choice of norm reflects the nature of the problem being studied. A global approximation considers the entire domain of the function and attempts to minimize the error everywhere. In contrast, a local approximation focuses on a neighborhood around a point, which is particularly useful for adaptive and meshless methods where local behavior is more important than global accuracy.

Common norms used in approximation theory include the maximum norm, which measures the largest deviation between the exact and approximate functions, and is often used in uniform approximation problems. The L2L^2L2-norm, commonly associated with least squares approximation, measures the mean square error and is widely used in engineering applications due to its mathematical convenience and physical interpretation. In some applications, weighted norms are employed to emphasize accuracy in specific regions of the domain, such as near boundaries or regions with steep gradients.

Approximation theory not only provides techniques for constructing approximations, but also establishes theoretical results related to error bounds**,** convergence rates**,** and stability. Error bounds quantify how close the approximation is to the true function, while convergence rates describe how the approximation improves as the number of basis functions or data points increases.

Page 104-114 © MANTECH PUBLICATIONS 2019. All Rights Reserved

Stability ensures that small perturbations in data do not lead to large errors in the approximation, which is essential for reliable numerical computation.

Polynomial Approximation

Polynomial approximation is one of the earliest and most fundamental areas of approximation theory. Polynomials are attractive because they are simple to evaluate, differentiate, and integrate. A classical result in this area is the Weierstrass Approximation Theorem, which states that any continuous function defined on a closed and bounded interval can be approximated uniformly by a polynomial to any desired degree of accuracy.

Several polynomial approximation techniques are commonly used. Taylor series approximation represents a function locally around a point using derivatives, and is particularly useful for smooth functions. However, Taylor series approximations are inherently local and may not perform well far from the expansion point.

Interpolation polynomials**,** such as those in Lagrange or Newton form, construct a polynomial that exactly matches the function values at a given set of points. These methods are straightforward and provide exact fitting at interpolation nodes. However, when high-degree polynomials are used over large intervals, interpolation may lead to large oscillations near the boundaries, a phenomenon known as Runge's phenomenon**.** This issue limits the practical use of high-order global polynomial interpolation.

To overcome these difficulties, Chebyshev polynomial approximation is often employed. Chebyshev polynomials minimize the maximum error and provide near-optimal approximations with improved numerical stability. By choosing interpolation points based on Chebyshev nodes rather than equally spaced points, oscillations can be significantly reduced.

Despite their limitations, polynomial approximations remain widely used due to their simplicity and efficiency. They also serve as building blocks for more advanced approximation schemes used in meshless methods.

Page 104-114 © MANTECH PUBLICATIONS 2019. All Rights Reserved

Spline Approximation

Spline approximation was developed to address the shortcomings of global polynomial approximation, particularly the lack of local control and numerical instability. Splines are piecewise polynomial functions defined over subintervals of the domain, with smoothness conditions imposed at the joining points, known as knots**.** These smoothness conditions ensure continuity of the function and its derivatives up to a specified order.

Among various spline types, cubic splines are the most commonly used because they offer a good balance between smoothness and computational efficiency. Cubic splines ensure continuity of the function as well as its first and second derivatives, which makes them suitable for problems involving smooth physical behavior.

One of the main advantages of spline approximation is its local nature. Modifying the function in one subinterval affects only a small region, unlike global polynomial approximation where changes influence the entire domain. This local control leads to improved numerical stability and makes splines particularly useful in problems with complex geometries or localized features.

In the context of meshless methods, spline-like behavior is often achieved through **local** approximation techniques such as the moving least squares (MLS) method**.** Although MLS does not explicitly construct splines, it produces smooth, locally supported approximations that resemble spline functions. This connection highlights the strong role of spline concepts within modern meshless numerical methods.

Scattered Data Approximation

Classical approximation and interpolation techniques are generally developed under the assumption that data points lie on structured or semi-structured grids. However, in many practical problems—especially those involving complex geometries, evolving domains, or higher dimensions—such structured data is difficult or expensive to generate. Meshless methods naturally operate on scattered sets of nodes, without requiring any predefined connectivity. As a result,

Page 104-114 © MANTECH PUBLICATIONS 2019. All Rights Reserved

scattered data approximation becomes a central and unavoidable component of meshless numerical techniques.

The main challenge in scattered data approximation lies in constructing stable and accurate approximating functions using only nodal positions and function values. Approximation theory provides several powerful tools to address this problem, among which radial basis functions and moving least squares methods are the most widely used. These techniques allow smooth and flexible approximations over irregularly distributed points and form the mathematical foundation of many meshless formulations.

Radial Basis Function (RBF) Approximation

Radial basis function approximation is one of the most popular methods for scattered data approximation due to its simplicity, generality, and strong theoretical foundation. In RBF methods, the approximating function is expressed as a linear combination of radially symmetric basis functions centered at the given data points. The general form of an RBF approximation is given by

s(x)=∑i=1Nλi ϕ(‖x−xi‖),s(x) = \sum_{i=1}^{N} \lambda_i \, \phi(\|x - x_i\|),s(x)=i=1∑Nλi ϕ(‖x−xi‖),

where ϕ\phiϕ is a radial basis function that depends only on the distance between the evaluation point xxx and the center xix_ixi, λi\lambda_iλi are unknown coefficients, and NNN is the total number of nodes. The coefficients are determined by enforcing interpolation or approximation conditions, often leading to a system of linear equations.

A key feature of RBFs is that they are dimension-independent, meaning the same formulation applies in one, two, or higher dimensions without significant modification. This makes RBF methods particularly attractive for high-dimensional problems and complex geometries.

Several types of radial basis functions are commonly used in practice. Gaussian RBFs offer excellent smoothness and accuracy but depend on a shape parameter that strongly influences numerical stability. Multiquadric and inverse multiquadric functions are widely used due to their

Page 104-114 © MANTECH PUBLICATIONS 2019. All Rights Reserved

good approximation properties and flexibility. Thin plate spline functions, on the other hand, are particularly useful for smooth surface fitting and do not require a shape parameter.

Despite their advantages, RBF methods also have certain limitations. The resulting system matrices are generally dense**,** leading to high memory requirements and computational cost, especially for large-scale problems. In addition, the choice of the shape parameter plays a crucial role in balancing accuracy and stability, and selecting an optimal value remains an active research topic. Nevertheless, RBF approximation remains one of the most powerful tools for meshless and scattered data problems.

Moving Least Squares (MLS) Approximation

The moving least squares (MLS) method is another fundamental technique for scattered data approximation and is extensively used in meshless methods. Unlike interpolation-based approaches, MLS constructs an approximation by minimizing a weighted least squares error over a local neighborhood around each evaluation point. This local nature makes MLS particularly suitable for problems involving irregular node distributions.

In MLS approximation, the function is approximated by a polynomial whose coefficients vary spatially. At each point of interest, a local approximation is obtained by minimizing the difference between the polynomial and the nodal data, weighted by a function that decreases with distance. As the point moves through the domain, the approximation “moves” with it, which explains the name moving least squares.

One of the major strengths of MLS approximation is its smoothness**.** The resulting approximating functions are generally continuous and possess higher-order derivatives, depending on the chosen polynomial basis. This property is especially important in meshless methods for solving partial differential equations, where smooth trial functions are required.

Page 104-114 © MANTECH PUBLICATIONS 2019. All Rights Reserved

MLS approximation is widely used to construct shape functions in meshless methods such as the Element-Free Galerkin method and the Meshless Local Petrov–Galerkin method. These shape functions offer great flexibility and can easily handle scattered nodes and irregular domains.

However, MLS also has some drawbacks. Since MLS does not satisfy the Kronecker delta property, the approximated function does not necessarily pass through the nodal values. This makes the imposition of essential (Dirichlet) boundary conditions more complicated compared to finite element methods. Various techniques, such as penalty methods and Lagrange multipliers, are often employed to address this issue.

Overall, moving least squares approximation provides a robust and flexible framework for scattered data approximation and continues to play a vital role in the development of modern meshless numerical methods.

Overview of Meshless Methods

Meshless methods, also known as meshfree methods, eliminate the need for element connectivity. Only a set of nodes and a support domain are required.

Classification of Meshless Methods

Meshless methods can be broadly classified as:

- Strong form methods,
- Weak form methods.

Strong form methods directly approximate derivatives in the governing equations, while weak form methods use variational formulations similar to FEM.

Table 1: Classification of Common Meshless Methods

Method Name	Approximation Technique	Formulation	Typical Applications
SPH	Kernel approximation	Strong form	Fluid dynamics
EFG	MLS	Weak form	Solid mechanics

Page 104-114 © MANTECH PUBLICATIONS 2019. All Rights Reserved

Method Name	Approximation Technique	Formulation	Typical Applications
MLPG	MLS / RBF	Local weak form	Heat transfer
RBF Collocation	RBF	Strong form	PDE solving

Element-Free Galerkin Method

The Element-Free Galerkin (EFG) method is one of the most popular meshless methods. It employs the Galerkin weak form and uses MLS approximation to construct trial and test functions. EFG is particularly suitable for problems involving large deformation and crack growth. However, numerical integration in EFG often requires background cells, which partially reduces its meshless nature.

Smoothed Particle Hydrodynamics

Smoothed Particle Hydrodynamics (SPH) is a particle-based meshless method originally developed for astrophysical simulations. It uses kernel functions to approximate field variables and their derivatives.

SPH has been successfully applied to fluid flow, free surface problems, and impact dynamics. Despite its advantages, SPH can suffer from consistency and stability issues, especially near boundaries.

Applications of Approximation Theory and Meshless Methods

Meshless methods supported by approximation theory have found applications in many areas:

- **Solid Mechanics:** crack propagation, fracture analysis.
- **Fluid Dynamics:** free surface flows, multiphase flows.
- **Heat Transfer:** problems with moving heat sources.
- **Biomechanics:** modeling of soft tissues.
- **Electromagnetics:** scattered field approximation.

Page 104-114 © MANTECH PUBLICATIONS 2019. All Rights Reserved

In mesh-based methods, the domain is divided into structured or unstructured elements. In meshless methods, only nodes are scattered throughout the domain, and approximation is constructed locally using support domains.

Advantages and Limitations

Advantages

- No need for mesh generation.
- Easy handling of large deformations.
- Flexibility in node distribution.
- Higher-order continuity possible.

Limitations

- Computational cost is often higher.
- Implementation is more complex.
- Boundary condition enforcement is difficult.
- Numerical integration may still need background cells.

Current Challenges and Future Directions

Although meshless methods have matured significantly, several challenges remain. Improving computational efficiency, developing robust boundary condition techniques, and ensuring numerical stability are active research areas. Hybrid approaches combining mesh-based and meshless methods are also gaining attention.

Future research may focus on adaptive node refinement, parallel implementations, and machine learning-assisted approximation techniques. The strong link between approximation theory and data-driven methods also opens new opportunities.

CONCLUSION

Approximation theory provides the essential mathematical framework for meshless methods. From polynomial and spline approximations to advanced radial basis and moving least squares techniques, approximation concepts enable the construction of flexible and accurate numerical

Page 104-114 © MANTECH PUBLICATIONS 2019. All Rights Reserved

schemes. Meshless methods, in turn, extend the applicability of numerical analysis to complex and evolving problems where traditional mesh-based approaches struggle.

This review has presented a detailed discussion on approximation theory and its role in meshless methods, along with practical applications and challenges. Although some issues still exist, the continued development of approximation techniques and computational resources suggests that meshless methods will play an increasingly important role in scientific and engineering computations.

REFERENCES

1. Cheney, E. W. (1982). *Introduction to Approximation Theory*. McGraw-Hill.
2. Powell, M. J. D. (1992). The theory of radial basis function approximation. *Advances in Numerical Analysis*, 2, 105–210.
3. Lancaster, P., & Salkauskas, K. (1986). *Curve and Surface Fitting*. Academic Press.
4. Belytschko, T., Lu, Y. Y., & Gu, L. (1994). Element-free Galerkin methods. *International Journal for Numerical Methods in Engineering*, 37, 229–256.
5. Liu, G. R. (2003). *Mesh Free Methods: Moving Beyond the Finite Element Method*. CRC Press.
6. Wendland, H. (2005). *Scattered Data Approximation*. Cambridge University Press.
7. Monaghan, J. J. (1992). Smoothed particle hydrodynamics. *Annual Review of Astronomy and Astrophysics*, 30, 543–574.
8. Atluri, S. N., & Zhu, T. (1998). A new meshless local Petrov–Galerkin approach. *Computational Mechanics*, 22, 117–127.
9. Fasshauer, G. E. (2007). *Meshfree Approximation Methods with MATLAB*. World Scientific.
10. Griebel, M., & Schweitzer, M. A. (2002). *Meshfree Methods for Partial Differential Equations*. Springer.

Page 104-114 © MANTECH PUBLICATIONS 2019. All Rights Reserved