

Homoeopathic Therapeutics in Dental Surgery and Perioperative Care: Adjunctive Management and Wound Healing

Dr. Meghana Patil¹, Dr. Irfan Sheikh², Tara Krishnan³, Amit Joshi⁴

ABSTRACT

Within homoeopathic practice, medicines from the materia medica are used for a range of minor dental complaints and as adjuncts in the perioperative period, the best known being Arnica montana for the soreness and bruising that follow injury or surgery. This paper reviews the role of homoeopathic therapeutics in dental and perioperative care, defining its adjunctive function, its application to wound healing and recovery, and its limitations. A review-based methodology is adopted, drawing on the homoeopathic materia medica and clinical literature, on standard oral-surgery references, and on the available evidence. The analysis indicates that homoeopathic use in this field is described chiefly for minor dental complaints, such as uncomplicated toothache, teething, and aphthous ulcers, and for adjunctive perioperative support across the pre-operative, post-operative, and wound-healing phases. A structured framework is presented comprising a perioperative timeline of adjunctive measures, a mapping of representative remedies to injury types, and a set of dental red flags requiring surgical or emergency care. The study emphasises strongly that homoeopathy is not a substitute for dental or surgical procedures, local anaesthesia, drainage of abscesses, antibiotics where indicated, or emergency care, and that conditions such as spreading infection, facial swelling, airway compromise, uncontrolled bleeding, and dental trauma require prompt conventional management. It is further noted that the clinical evidence is limited and mixed, with systematic reviews of widely studied agents such as homoeopathic arnica finding the evidence unconvincing. The paper concludes that any role for homoeopathic therapeutics here is confined to minor complaints and adjunctive support, must operate within clear referral safeguards, and must never delay necessary surgical, anaesthetic, or antibiotic treatment.

KEYWORDS: *Homoeopathic Therapeutics, Dental Surgery, Perioperative Care, Arnica, Wound Healing, Adjunctive Treatment, Referral, Materia Medica*

INTRODUCTION

Dental and oral-surgical care encompasses procedures ranging from simple restorations and extractions to the management of infection, trauma, and complex maxillofacial surgery, alongside the routine perioperative concerns of anxiety, pain, bleeding, bruising, and wound healing [7]. Within the homoeopathic system, medicines from the materia medica have long been described for a number of minor dental complaints and, most prominently, as adjuncts in the perioperative period; the use of Arnica montana for the soreness and bruising that follow injury or operation is the most widely known example and extends across surgical practice generally [1], [4], [5].

The central problem this paper addresses is how to define the legitimate scope of homoeopathic therapeutics in dental and perioperative care responsibly. The relevant questions are which minor complaints might reasonably be managed or supported with homoeopathic medicines, how such measures should function as adjuncts to conventional dental and surgical care, and, crucially, where the firm boundaries lie beyond which surgical, anaesthetic, or antibiotic treatment is mandatory and must not be delayed. As in other surgical fields, some dental and oral conditions are urgent: a spreading dental infection, a swelling that threatens the airway, uncontrolled bleeding, or significant trauma all require prompt conventional management, and failure to provide it can lead to serious harm.

A responsible account must therefore pair any description of homoeopathic adjuncts with explicit recognition of conventional priorities and referral safeguards, and with an honest statement of the evidence. Accordingly, this paper reviews the homoeopathic medicines described for dental complaints and perioperative support, presents a framework covering the perioperative period, wound healing, and dental red flags, and emphasises throughout that homoeopathy is an adjunct at most, never a substitute for dental and surgical procedures, anaesthesia, drainage, antibiotics, or emergency care, and that the supporting evidence is limited and mixed [5], [9], [10].

LITERATURE REVIEW

The homoeopathic materia medica and works of therapeutics describe medicines used for several minor dental and oral complaints, selected according to the individual symptom picture [1], [2], [3]. Commonly described medicines include Chamomilla for the unbearable toothache and irritability associated with dentition, particularly in children, and Plantago and Belladonna for acute toothache with inflammation; Mercurius solubilis and Hepar sulphuris where oral suppuration is present; Borax and Mercurius for aphthous ulceration and stomatitis; and Calcareo phosphorica in the context of dentition. For procedural anxiety, medicines such as Gelsemium, Aconite, and Argentum nitricum are described. These accounts represent homoeopathic prescribing tradition rather than established clinical efficacy, and they pertain to minor complaints [1], [4].

A larger and more distinctive strand concerns adjunctive perioperative use and wound healing across surgical practice, including dental and oral surgery. Here the medicines most frequently described are Arnica montana for the bruising, soreness, and tendency to bleeding that follow trauma and operation; Staphysagria for clean incised surgical wounds; Calendula officinalis as a wound-healing and antiseptic application, including as a mouth rinse; Hypericum perforatum for injuries to nerve-rich tissues and associated pain; Ledum palustre for puncture-type wounds; Bellis perennis for deeper tissue surgery; and Phosphorus for post-anaesthetic nausea and a bleeding tendency [1], [4], [6]. These are described as adjuncts intended to support comfort and recovery alongside conventional care. The standard oral-surgery literature, by contrast, defines the procedures, anaesthesia, drainage, and antibiotic therapy that constitute the primary management of dental disease, infection, and trauma, and specifies the emergencies that require immediate attention [7].

The clinical evidence for homoeopathy in this field is limited and mixed. Homoeopathic arnica, the most studied agent, has been examined in numerous placebo-controlled trials around surgery, including dental and orthopaedic procedures, with inconsistent results; systematic reviews have concluded that the evidence does not convincingly demonstrate efficacy beyond placebo [5], [6]. Authoritative overviews of the broader homoeopathic evidence base have likewise found no convincing support for efficacy across health conditions [9], [10]. This evidential context is essential to any responsible discussion. Critically, abscesses and spreading infections require drainage and, where indicated, antibiotics; surgical conditions require

surgery; and none of these can be safely managed by homoeopathic medicines alone. The literature therefore supports, at most, a bounded adjunctive role operating strictly alongside and subordinate to conventional dental and surgical care.

RESEARCH GAP

Descriptions of homoeopathic medicines for dental complaints and perioperative support are scattered across the materia medica and therapeutics literature and are often presented without a clear, structured account of their adjunctive role, their place within the perioperative period, and the boundaries that require surgical or emergency care. Conventional dental and oral-surgical priorities, in turn, are defined in specialty texts but are rarely integrated with the homoeopathic literature. There is therefore a need for a consolidated framework that defines the safe, adjunctive scope of homoeopathic therapeutics in dental and perioperative care, maps it onto the phases of treatment and to wound healing, embeds it within explicit referral safeguards, and presents the evidential limitations honestly.

OBJECTIVES

The specific objectives of this study are:

1. To review the homoeopathic medicines described for minor dental complaints and perioperative support.
2. To define the adjunctive scope of homoeopathic therapeutics across the perioperative period and in wound healing.
3. To present dental red-flag criteria requiring surgical or emergency care.
4. To state honestly the limitations and the evidential status of homoeopathy in this field.
5. To identify directions for safe, integrative practice and future research.

METHODOLOGY

A review-based methodology was adopted. The homoeopathic medicines described for dental complaints and perioperative support were compiled from the materia medica and works of homoeopathic therapeutics; the conventional priorities and emergency criteria were drawn from standard oral-surgery references; and the evidential status was characterised from systematic reviews and national assessments [1]-[10]. The information was organised into a framework comprising a perioperative timeline, a mapping of representative remedies to injury types, and a set of dental red flags.

Perioperative Framework

The adjunctive use of homoeopathic medicines across the perioperative period is represented in Figure 1 as a timeline through the pre-operative, operative, early post-operative, and wound-healing phases. In the pre-operative phase, medicines such as Gelsemium and Aconite are described for anxiety; the operative phase is conducted under conventional anaesthesia and surgery, which homoeopathy does not replace; the early post-operative phase concerns trauma, bruising, and pain, for which Arnica and Hypericum are described; and the wound-healing phase concerns recovery, for which Calendula and Staphysagria are described, with Phosphorus mentioned for post-anaesthetic nausea or bleeding tendency. The figure states explicitly that conventional anaesthesia, surgery, analgesia, and antibiotics remain the primary treatment [6], [7].




Figure 1: Adjunctive homoeopathic measures across the perioperative period, alongside primary conventional care

Wound Mapping and Dental Red Flags

Representative trauma and wound remedies are mapped to injury types in Figure 2, associating, for example, blunt trauma and bruising with Arnica, clean incised wounds with Staphysagria, open wounds and antisepsis with Calendula, nerve-rich injuries with Hypericum, and puncture wounds with Ledum. This mapping reflects homoeopathic prescribing tradition for adjunctive wound care. The dental conditions and emergencies requiring conventional surgical or urgent management are organised in Figure 3 and Table 3, ensuring that serious presentations are identified and referred rather than managed conservatively [7].




Figure 2: Representative homoeopathic trauma and wound remedies mapped to injury types (adjunctive use)

RESULTS AND FINDINGS

Medicines Described for Dental Complaints and Perioperative Support

Table 1 summarises the homoeopathic medicines commonly described for minor dental complaints, and Table 2 summarises those described across the phases of perioperative care. The findings show that described use concentrates on minor, largely self-limiting complaints, such as uncomplicated toothache, teething, and mouth ulcers, and on adjunctive support around procedures, the situations in which conservative or supportive measures are generally reasonable alongside standard care. The medicines are selected on the individual symptom picture in accordance with homoeopathic principles. It must be emphasised that these tables describe prescribing tradition, not demonstrated efficacy, and that they apply only to minor presentations and to adjunctive use; none is appropriate as a substitute for dental procedures, drainage, anaesthesia, antibiotics, or surgery [1], [5], [9].

Table 1: Minor dental complaints and representative homoeopathic medicines (adjunctive / minor presentations only).

Complaint	Typical homoeopathic indication	Representative medicines
Toothache (uncomplicated)	Acute pain with inflammation	Belladonna, Plantago, Chamomilla
Teething (dentition)	Painful, irritable dentition	Chamomilla, Calcarea phos

Complaint	Typical homoeopathic indication	Representative medicines
Aphthous ulcers / stomatitis	Oral mucosal soreness	Borax, Mercurius solubilis
Post-procedure soreness	Bruising, soreness (adjunct)	Arnica, Staphysagria
Procedural anxiety	Apprehension before treatment	Gelsemium, Aconite, Argentum nit.

Table 2: Perioperative phases and representative adjunctive homoeopathic measures.

Phase	Adjunctive concern	Representative medicines
Pre-operative	Anxiety, apprehension	Gelsemium, Aconite, Argentum nit.
Operative	Conventional anaesthesia & surgery	(not replaced by homoeopathy)
Early post-operative	Trauma, bruising, pain	Arnica, Hypericum
Wound healing	Recovery, antiseptis	Calendula, Staphysagria
Post-anaesthetic	Nausea, bleeding tendency	Phosphorus

Boundaries and Dental Red Flags

The analysis identifies a clear, non-negotiable boundary between minor complaints amenable to conservative or adjunctive measures and conditions requiring conventional surgical, anaesthetic, or antibiotic management. Figure 3 and Table 3 set out the dental red flags that mandate prompt care, including facial or neck swelling, spreading infection or fever, difficulty breathing or swallowing, uncontrolled bleeding, dental trauma or an avulsed tooth, and abscess or severe unremitting pain. The finding is that recognition of these features and immediate appropriate management, including drainage, antibiotics, anaesthesia, or surgery as indicated, is the most important element of safe practice and overrides any consideration of homoeopathic treatment [7]. Adjunctive homoeopathic measures, where used at all, are confined to minor complaints and to supportive roles around otherwise properly managed procedures.




Figure 3: Dental red-flag features requiring surgical, antibiotic, or emergency care.

Table 3: Dental red flags and required conventional management.

Red-flag feature	Required conventional management
Facial / neck swelling; spreading infection	Drainage and/or antibiotics; urgent review
Difficulty breathing or swallowing	Emergency care (airway risk)
Uncontrolled bleeding	Haemostatic and surgical management
Dental trauma / avulsed tooth	Urgent dental / surgical care
Abscess; severe unremitting pain	Drainage; definitive dental treatment

DISCUSSION

The central message of this study is that homoeopathic therapeutics in dental and perioperative care can be considered only as an adjunct, tightly bounded and subordinate to conventional treatment. The perioperative framework (Figure 1) positions any homoeopathic measure alongside, and explicitly after, conventional anaesthesia, surgery, analgesia, and antibiotics, which remain primary at every stage. The wound mapping (Figure 2) reflects homoeopathic prescribing tradition for supportive recovery, not a claim to replace surgical wound

management. The dental red flags (Figure 3) ensure that infection, trauma, and airway-threatening or bleeding emergencies are identified and managed conventionally without delay. This structure mirrors safe surgical practice and is the only responsible context for the topic [6], [7].

The most important practical implication is that homoeopathy must never delay or replace necessary dental and surgical treatment. Dental infections can spread rapidly and, in severe cases, threaten the airway or become systemic; abscesses require drainage and frequently antibiotics; trauma and uncontrolled bleeding require prompt intervention. An approach that substituted homoeopathic medicines for these treatments would risk serious harm. Even where adjunctive measures such as Arnica are used after a properly performed procedure, they should be understood as supportive additions for comfort, used alongside standard analgesia and aftercare rather than instead of them, and discontinued in favour of review if recovery does not proceed normally [5], [7].

An honest account of the evidence is essential. Homoeopathic arnica, despite being among the most studied homoeopathic interventions in surgery, has not been shown by systematic review to be convincingly effective beyond placebo, and the broader evidence base for homoeopathy has been judged unconvincing by authoritative assessments [5], [9], [10]. Patients undergoing dental or surgical procedures are best served by transparent information about this uncertainty and by clear prioritisation of treatments of established benefit. The appropriate stance is therefore cautious and integrative: any homoeopathic use confined to minor complaints and adjunctive support, conventional care always primary, referral safeguards rigorously maintained, and evidential limitations communicated openly. Within these constraints, the contribution of homoeopathic therapeutics in dental and perioperative care is, at most, a limited adjunctive one.

LIMITATIONS

This study is a review and framework synthesis rather than a clinical trial, and it does not provide evidence of efficacy for homoeopathic medicines in any dental or perioperative condition; the tables describe prescribing tradition, not demonstrated outcomes. The scope is limited to minor complaints and adjunctive support and to the question of safe use, and it does not address the detailed management of dental disease, infection, or trauma, which belongs to

dental and oral-surgical practice. The red-flag criteria are presented in summary form for orientation and do not replace clinical assessment by a qualified dental or medical practitioner. Because the evidence base is limited and mixed, the conclusions emphasise safe scope, conventional priority, and referral rather than therapeutic recommendation, and should be read as guidance toward caution and integration rather than endorsement of any specific treatment.

FUTURE SCOPE

Future work in this area can usefully proceed as follows:

- Development of clear guidance defining the safe, adjunctive scope of homoeopathic measures in dental and perioperative care.
- Integration of dental red-flag and referral protocols into any homoeopathic practice in this field.
- Rigorous, well-controlled trials of specific adjunctive uses, such as post-operative comfort, with appropriate outcome measures.
- Education emphasising recognition of dental infection, trauma, and airway emergencies and timely conventional management.
- Models of integrative dental care that keep conventional treatment primary and maintain clear referral pathways.
- Transparent communication of the evidential limitations to patients undergoing dental and surgical procedures.

Pursuing these directions would help ensure that any use of homoeopathic therapeutics in dental and perioperative care remains safe, appropriately bounded, and subordinate to necessary conventional treatment.

CONCLUSION

This paper has reviewed the role of homoeopathic therapeutics in dental and perioperative care and defined its adjunctive function, its application to wound healing, and its limitations. The analysis shows that homoeopathic use here is described principally for minor dental complaints, such as uncomplicated toothache, teething, and mouth ulcers, and for adjunctive support across the perioperative period, with *Arnica montana* for post-procedure bruising and soreness being the best-known example. A structured framework comprising a perioperative timeline, a wound-and-injury remedy mapping, and dental red-flag criteria defines the safe

limits of such use. The overriding conclusions are that homoeopathy is not a substitute for dental or surgical procedures, anaesthesia, drainage of abscesses, antibiotics, or emergency care; that red-flag conditions such as spreading infection, airway compromise, uncontrolled bleeding, and trauma require prompt conventional management; and that the clinical evidence is limited and mixed, with systematic reviews of homoeopathic arnica finding it unconvincing. Any role for homoeopathic therapeutics in this field must therefore be confined to minor complaints and adjunctive support, operate within clear referral safeguards, and never delay necessary surgical, anaesthetic, or antibiotic treatment.

REFERENCES

1. W. Boericke, *Pocket Manual of Homoeopathic Materia Medica with Repertory*. New Delhi, India: B. Jain Publishers.
2. J. T. Kent, *Repertory of the Homoeopathic Materia Medica*. New Delhi, India: B. Jain Publishers.
3. H. C. Allen, *Keynotes and Characteristics with Comparisons of Some of the Leading Remedies*. New Delhi, India: B. Jain Publishers.
4. E. B. Nash, *Leaders in Homoeopathic Therapeutics*. New Delhi, India: B. Jain Publishers.
5. E. Ernst and M. H. Pittler, "Efficacy of homeopathic arnica: A systematic review of placebo-controlled clinical trials," *Arch. Surg.*, vol. 133, no. 11, pp. 1187–1190, 1998.
6. B. Brinkhaus *et al.*, "Homeopathic arnica therapy in patients receiving knee surgery: Results of three randomised double-blind trials," *Complement. Ther. Med.*, vol. 14, no. 4, pp. 237–246, 2006.
7. J. R. Hupp, E. Ellis, and M. R. Tucker, *Contemporary Oral and Maxillofacial Surgery*, 7th ed. St. Louis, MO, USA: Elsevier, 2019.
8. S. Hahnemann, *Organon of Medicine*, 6th ed., W. Boericke, Trans. New Delhi, India: B. Jain Publishers.
9. E. Ernst, "A systematic review of systematic reviews of homeopathy," *Br. J. Clin. Pharmacol.*, vol. 54, no. 6, pp. 577–582, 2002.
10. National Health and Medical Research Council, *Evidence on the Effectiveness of Homeopathy for Treating Health Conditions*. Canberra, Australia: NHMRC, 2015.
11. World Health Organization, *Safety Issues in the Preparation of Homeopathic Medicines*. Geneva, Switzerland: WHO, 2009.

12. R. Murphy, *Homeopathic Medical Repertory*. New Delhi, India: B. Jain Publishers.
13. A. Shang *et al.*, “Are the clinical effects of homoeopathy placebo effects? Comparative study of placebo-controlled trials of homoeopathy and allopathy,” *Lancet*, vol. 366, no. 9487, pp. 726–732, 2005.
14. C. Lokken *et al.*, “Effect of homoeopathy on pain and other events after acute trauma: Placebo controlled trial with bilateral oral surgery,” *BMJ*, vol. 310, no. 6992, pp. 1439–1442, 1995.
15. D. M. Eisenberg *et al.*, “Complementary and alternative medicine use: Implications for clinical and dental practice,” *J. Am. Dent. Assoc.*, vol. 138, no. 1, pp. 95–101, 2007.
16. K. Linde *et al.*, “Are the clinical effects of homeopathy placebo effects? A meta-analysis of placebo-controlled trials,” *Lancet*, vol. 350, no. 9081, pp. 834–843, 1997.
17. Ministry of AYUSH, *Standard Treatment Guidelines for Homoeopathy*. New Delhi, India: Government of India.
18. S. B. Kayne, *Homeopathic Pharmacy: Theory and Practice*, 2nd ed. Edinburgh, U.K.: Elsevier Churchill Livingstone, 2006.

***Author for Correspondence**

Dr. Meghana Patil

E-mail: meghana.patil239@rediffmail.com

¹Lecturer, Department of Homoeopathic Therapeutics, J.R. Kissan Homoeopathic Medical College & Hospital

²Professor, Department of Homoeopathic Therapeutics, J.R. Kissan Homoeopathic Medical College & Hospital

^{3,4}PG Scholar, Department of Homoeopathic Therapeutics, J.R. Kissan Homoeopathic Medical College & Hospital

Received Date: March 05, 2026,

Accepted Date: March 19, 2026,

Published Date: March 26, 2026

Citation: Dr. Meghana Patil, Dr. Irfan Sheikh, Tara Krishnan, Amit Joshi. Homoeopathic Therapeutics in Dental Surgery and Perioperative Care: Adjunctive Management and Wound Healing. *Homoeopathic Surgical Innovations: Journal of ENT, Ophthalmology, Dental, and Therapeutics*. 2026; 3(1): 28-39p.