

Artificial Intelligence and Explainability in Earth Observation: Enhancing Transparency, Trust, and Decision-Making in Remote Sensing Applications

Dr. Kavita Nair¹, Arjun Deshmukh², Mayank Rawat³

Associate Professor¹, Research Scholars^{2,3}

Department of Civil Engineering¹, Department of CSE^{2,3}

College of Engineering, Pune (COEP)

Email ID: *kavitanai4033@gmail.com¹, arjun_deshmukh@yahoo.co.in²*

ABSTRACT

Artificial Intelligence (AI) has emerged as a transformative technology in Earth Observation (EO), enabling automated analysis of vast datasets obtained from satellites, drones, and sensors. These AI-driven systems contribute significantly to environmental monitoring, disaster management, urban planning, and climate modeling. However, the rapid adoption of AI in EO introduces a critical challenge—explainability. Explainable AI (XAI) ensures that model outputs are interpretable, trustworthy, and scientifically valid, especially when decisions impact environmental policy, safety, and sustainability. This paper explores the convergence of AI and explainability in Earth Observation, emphasizing methodologies, challenges, and future directions for transparent and interpretable models. The study provides a comprehensive view of the evolution, techniques, applications, and prospects of XAI in the EO domain.

KEYWORDS: *Artificial Intelligence, Explainable AI, Earth Observation, Remote Sensing, Interpretability, Machine Learning, Deep Learning, Environmental Monitoring, Transparency, Trust.*

INTRODUCTION

Earth Observation has entered an era of data abundance, driven by advanced satellite constellations, high-resolution sensors, and Internet of Things (IoT)-enabled ground systems. These technologies generate petabytes of data daily, capturing dynamic changes on Earth's

surface. Traditional data analysis methods are inadequate to handle this data scale, leading to the integration of AI-based systems such as machine learning (ML) and deep learning (DL).

AI models can automatically detect land-use changes, monitor deforestation, identify natural hazards, and estimate crop yields. However, most of these systems are black-box models—they provide predictions without explaining the underlying reasoning. For scientists, policymakers, and the general public, understanding how AI makes decisions is crucial for accountability and trust. Therefore, Explainable AI (XAI) has emerged as a bridge between complex AI algorithms and human interpretability in EO applications.

LITERATURE REVIEW

AI in Earth Observation:

AI has revolutionized EO data processing by automating feature extraction, classification, and anomaly detection. Convolutional Neural Networks (CNNs), Recurrent Neural Networks (RNNs), and Vision Transformers (ViTs) have demonstrated superior performance in recognizing spatial and temporal patterns. Studies have shown that AI-based approaches outperform traditional pixel-based classifiers such as Maximum Likelihood and Support Vector Machines. For instance, deep learning techniques have achieved remarkable accuracy in land cover mapping, urban sprawl detection, and flood extent analysis.

Explainability in AI:

Despite their success, deep learning models are often criticized for their “black-box” nature. Explainable AI (XAI) aims to make these models transparent and interpretable by providing insights into how decisions are made. In EO, interpretability is essential not only for scientific validation but also for policy formulation and ethical decision-making. Techniques like Layer-wise Relevance Propagation (LRP), SHapley Additive exPlanations (SHAP), and Local Interpretable Model-agnostic Explanations (LIME) have been increasingly used to visualize which features or regions contribute most to a model’s prediction.

Integration of AI and XAI in EO:

Recent research highlights the integration of XAI tools within EO pipelines. For example, using SHAP in satellite-based vegetation health prediction allows scientists to understand which spectral indices contribute to vegetation stress. Similarly, attention mechanisms in deep neural

networks can highlight crucial spatial regions for wildfire detection or glacier retreat analysis. Thus, XAI serves as a scientific audit mechanism that enhances confidence in AI-driven EO models.

METHODOLOGIES IN AI AND EXPLAINABILITY

Table 1: Comparison of AI and XAI Techniques in Earth Observation

Technique	Category	Application Area in EO	Explainability Feature	Advantages	Limitations
CNN (Convolutional Neural Network)	Deep Learning	Land cover classification, object detection	Visualization through Grad-CAM	High spatial accuracy	Low interpretability
Random Forest	Machine Learning	Soil and vegetation mapping	Feature importance ranking	Easy to interpret	Limited for complex data
SHAP (SHapley Additive ExPlanations)	Post-hoc XAI	Crop yield prediction, drought monitoring	Attribute-level explanation	Model-agnostic	High computation cost
LIME (Local Interpretable Model-agnostic Explanations)	Model-Agnostic XAI	Disaster assessment	Localized feature interpretation	Human-readable	Inconsistent explanations
Attention-based Transformers	Deep Learning	Change detection, image segmentation	Attention heatmaps	High performance, interpretable focus	Computationally intensive




Figure 1: Workflow of Explainable AI in Earth Observation

1. Machine Learning and Deep Learning Models:

AI models for EO typically involve supervised and unsupervised learning. Supervised models such as Random Forests and Support Vector Machines classify satellite images using labeled datasets. Deep Learning models like CNNs and U-Net architectures are used for semantic segmentation tasks such as identifying water bodies, vegetation, and built-up areas.

2. Explainable AI Techniques:

- **Feature Importance Methods:** SHAP and LIME assign contribution scores to input features, allowing users to see which spectral bands or indices influenced the classification.
- **Visualization-based Explainability:** Grad-CAM and saliency maps visualize image regions that strongly affect model predictions.
- **Model Simplification:** Decision trees or surrogate models are trained to mimic complex models while remaining interpretable.
- **Concept-based Explanations:** These connect model outputs with human-understandable concepts, such as vegetation index or soil moisture.

3. Hybrid Approaches:

Researchers are developing hybrid frameworks combining physical models with data-driven AI. Such models retain interpretability while leveraging AI's predictive power. For example, combining hydrological models with deep learning improves flood forecasting accuracy while maintaining transparency.

APPLICATIONS OF XAI IN EARTH OBSERVATION

Table 2: Applications of Explainable AI in Earth Observation Domains

Application Domain	Objective	AI Technique Used	Explainability Tool	Outcome/Benefit
Climate Change Monitoring	Track glacier retreat, carbon emissions	CNN, Autoencoder	SHAP, LRP	Transparent climate trend insights
Precision Agriculture	Crop health & yield estimation	Random Forest, LSTM	LIME, SHAP	Clear identification of stress indicators

Application Domain	Objective	AI Technique Used	Explainability Tool	Outcome/Benefit
Disaster Management	Flood and wildfire detection	U-Net, CNN	Grad-CAM	Visual interpretability of disaster zones
Urban Planning	Land use & change detection	Vision Transformer	Attention Maps	Transparent urban growth insights
Marine and Coastal Studies	Pollution & coral bleaching tracking	RNN, CNN	Saliency Maps	Explainable marine ecosystem changes

1. Climate Change and Environmental Monitoring

Climate change is one of the most pressing global issues, and remote sensing satellites continuously capture vast data related to temperature variations, atmospheric composition, and land-use changes. However, traditional AI models that analyze such data often lack interpretability, making it difficult for scientists to understand *why* a model predicts a particular environmental change.

Explainable AI resolves this challenge by visualizing and quantifying how different environmental parameters—such as surface temperature, vegetation index, or carbon concentration—contribute to the overall prediction. For instance, SHAP (SHapley Additive exPlanations) can quantify which climatic variables most influence forest degradation or glacial retreat. Similarly, Grad-CAM heatmaps can highlight satellite image regions that show the most significant evidence of melting ice or deforestation.

By identifying causal relationships between variables like greenhouse gas emissions, deforestation rates, and sea-level rise, XAI enables scientists to generate more interpretable climate models. These models support the creation of data-driven environmental policies and promote transparency in international climate agreements and sustainability programs.

2. Agriculture and Crop Monitoring

The agriculture sector benefits immensely from Explainable AI-based EO systems, especially

in precision agriculture, which requires continuous monitoring of crop health, soil quality, and moisture levels. Traditional deep learning models can predict crop yield or detect pest infestations with high accuracy, but they often fail to clarify which features—such as spectral bands, soil temperature, or rainfall—drive their predictions.

Using XAI techniques like LIME (Local Interpretable Model-Agnostic Explanations) and SHAP, agronomists can determine the relative influence of various environmental variables on crop conditions. For example, SHAP can reveal whether Normalized Difference Vegetation Index (NDVI) or soil moisture content has a stronger correlation with reduced yield. Such explainable insights enable data-backed decisions on irrigation scheduling, fertilizer application, and pest control strategies.

Moreover, explainable AI supports transparent crop insurance assessments by helping stakeholders understand why certain regions are classified as drought-prone or high-risk. This transparency enhances trust among farmers, policymakers, and financial institutions, promoting equitable agricultural practices.

3. Disaster Management

Explainable AI has become indispensable in disaster risk prediction and post-event analysis. During natural calamities like floods, earthquakes, or wildfires, time-critical decision-making depends heavily on AI-driven models that analyze satellite data in real time. However, black-box predictions may not always be reliable for life-saving operations.

By incorporating explainability, AI models can highlight key risk indicators that trigger alerts. For instance, saliency maps can identify flood-prone areas by highlighting low-elevation zones or high soil saturation regions. In wildfire prediction, attention-based networks can pinpoint vegetation density and wind patterns as contributing factors to fire spread.

XAI improves accountability and transparency in emergency response by allowing disaster management authorities to understand why specific regions are classified as high-risk. It also enables post-disaster damage assessment by clearly visualizing affected zones, optimizing resource allocation, and supporting recovery planning.

4. Urban and Land Use Planning

Urbanization is accelerating globally, leading to rapid land cover transformation and ecosystem disruption. Earth Observation data combined with Explainable AI assists urban planners in analyzing spatial growth patterns, green cover loss, and infrastructure expansion with higher clarity.

Through explainability techniques, planners can understand which environmental and socio-economic features influence model outputs. For example, SHAP can identify that proximity to major roads and population density are the strongest predictors of urban sprawl. Similarly, attention maps in Vision Transformer (ViT) models highlight areas with rapid land-use conversion, such as farmland turning into residential zones.

By integrating XAI-driven insights, policymakers can develop sustainable urban development strategies, ensuring a balance between modernization and ecological preservation. This transparency also supports public trust and community engagement in urban decision-making processes.

5. Ocean and Coastal Studies

Coastal regions and marine ecosystems are highly dynamic and vulnerable to climate change, pollution, and human activities. Explainable AI techniques enable researchers to monitor and interpret changes in these environments with greater confidence.

For example, CNN-based ocean monitoring systems, when combined with Grad-CAM visualization, can pinpoint exact regions of oil spills, sediment deposition, or coral bleaching. Similarly, SHAP-based analyses help scientists understand the relative impact of sea surface temperature, chlorophyll concentration, and wave intensity on marine ecosystem degradation. These explainable insights facilitate international collaboration for marine conservation and policy transparency in environmental governance. Furthermore, coastal management agencies can use XAI to predict and mitigate erosion risks or tsunami hazards while understanding the rationale behind each forecasted event.

Thus, XAI serves as an essential analytical companion to traditional marine observation systems, enhancing both interpretability and environmental accountability.

CHALLENGES AND LIMITATIONS

1. Black-Box Nature of Deep Models:

Although XAI tools provide partial interpretability, complete transparency is still challenging. Deep models often involve millions of parameters, making human comprehension difficult.

2. Lack of Standardized Evaluation Metrics:

There is no universally accepted metric for evaluating the quality of explanations. What is interpretable to one user may not be understandable to another.

3. Data Quality and Heterogeneity:

EO data are collected from multiple sensors with varying resolutions and noise levels. These inconsistencies affect model reliability and interpretation.

4. Computational Complexity:

Implementing XAI increases computational cost and time, particularly when analyzing large-scale EO datasets.

5. Ethical and Policy Concerns:

Misinterpretation of AI decisions could lead to policy biases or incorrect environmental assessments. Ethical guidelines for explainability are still evolving.

SCOPE AND FUTURE PROSPECTS

1. Integration of Physical and Data-Driven Models:

The future of EO lies in combining AI-based approaches with physics-informed models to ensure both accuracy and interpretability.

2. Real-Time Explainability:

Developing real-time XAI systems that provide on-the-fly interpretations during disaster events can revolutionize decision support systems.

3. Federated and Edge Learning in EO:

Distributed AI models with explainability features can ensure data privacy while enabling large-scale collaboration across satellite networks.

4. Interdisciplinary Collaboration:

Bringing together experts from AI, remote sensing, geoscience, and ethics will foster the development of trustworthy AI systems in EO.

5. Policy and Governance:

Explainable AI will play a pivotal role in environmental governance, where transparency and accountability are paramount for decision-making.

RESULTS AND DISCUSSION



Figure 2: Grad-CAM Example Showing Explainable Land Cover Classification

Explainable AI has shown promising results in increasing stakeholder trust and improving scientific understanding. For instance, models using Grad-CAM for land cover classification demonstrate that the AI system focuses on the same regions as human analysts, validating its reasoning. Similarly, SHAP-based models for drought monitoring have revealed that vegetation indices and temperature anomalies are the most influential factors in model predictions.

The inclusion of explainability mechanisms not only improves user confidence but also enhances model generalization. By understanding how models interpret data, researchers can identify biases, improve data preprocessing steps, and refine model architectures.

CONCLUSION

Artificial Intelligence and Explainability together mark a transformative step in the evolution of Earth Observation. While AI accelerates data processing and predictive analysis,

explainability ensures that such automation remains transparent, interpretable, and ethically grounded. In environmental decision-making, trust is non-negotiable—hence, XAI provides the necessary framework to ensure that AI models are not only accurate but also accountable.

The journey ahead involves the integration of human-centered AI principles, where models complement human expertise rather than replace it. As Earth Observation continues to generate vast and complex data, the synergy between AI and explainability will be central to achieving a sustainable, transparent, and data-driven future for our planet.

REFERENCES

1. Adadi, A., & Berrada, M. (2018). Peeking inside the black-box: A survey on Explainable Artificial Intelligence (XAI). *IEEE Access*, 6, 52138–52160. <https://doi.org/10.1109/ACCESS.2018.2870052>
2. Alam, F., Mehmood, R., Katib, I., & Albeshri, A. (2020). Analysis of eight data mining algorithms for smarter Internet of Things (IoT). *IEEE Access*, 8, 110548–110563. <https://doi.org/10.1109/ACCESS.2020.2996938>
3. Anderson, K., & Gaston, K. J. (2013). Lightweight unmanned aerial vehicles will revolutionize spatial ecology. *Frontiers in Ecology and the Environment*, 11(3), 138–146.
4. Barredo Arrieta, A., Díaz-Rodríguez, N., Ser, J. D., Bennetot, A., Tabik, S., Barbado, A., ... & Herrera, F. (2020). Explainable Artificial Intelligence (XAI): Concepts, taxonomies, opportunities and challenges toward responsible AI. *Information Fusion*, 58, 82–115. <https://doi.org/10.1016/j.inffus.2019.12.012>
5. Bastin, L., & Buchanan, G. (2015). Spatial data infrastructures for environmental management. *Environmental Modelling & Software*, 68, 202–214.
6. Camps-Valls, G., Sejdinovic, D., Runge, J., & Reichstein, M. (2019). A perspective on Gaussian processes for Earth observation. *Nature*, 568(7753), 370–378.
7. Cheng, G., Yang, C., & Han, J. (2020). Learning with weak supervision for geospatial object detection and land-cover classification. *ISPRS Journal of Photogrammetry and Remote Sensing*, 167, 124–137.
8. Coppola, A. I., & Roderick, M. L. (2021). The role of explainable AI in understanding satellite-based climate indicators. *Remote Sensing of Environment*, 265, 112642.
9. Drusch, M., et al. (2012). Sentinel-2: ESA's optical high-resolution mission for GMES

operational services. *Remote Sensing of Environment*, 120, 25–36.

10. Ehsani, M. R., & Pradhan, B. (2021). Explainable deep learning for landslide susceptibility mapping using SHAP and LIME. *Geoscience Frontiers*, 12(6), 101224.