MANTECH Publications
Journal of Remote Sensing, Environmental Science & Geotechnical Engineering
Volume 8, Issue 1, January-April, 2023

Advances in Remote Sensing Techniques for Monitoring Greenhouse Gases and Their Sources and Sinks

Lokesh J K, Vikas Singh Rajput
Assistant Professor[1], Student[2]
Department of Civil Engineering
NMAM Institute of Technology Nitte, Karnataka
Corresponding Authors' Email: - *vikashrajput5@gmail.com*

Abstract

The increase in atmospheric greenhouse gases, particularly carbon dioxide, has led to global climate change. Remote sensing has become an essential tool for monitoring greenhouse gases and their sources and sinks on a global scale. This paper presents a review of the current state of remote sensing of greenhouse gases and their sources and sinks. The paper focuses on carbon dioxide (CO2), methane (CH4), and nitrous oxide (N2O), which are the three most significant greenhouse gases responsible for climate change. The paper describes the remote sensing techniques used to measure these gases and their sources and sinks and the challenges faced by these techniques. The paper also discusses the future prospects of remote sensing in monitoring greenhouse gases and their sources and sinks.

Keywords: *Remote sensing, greenhouse gases, sources, sinks, carbon dioxide, methane, nitrous oxide, satellite, ground-based, airborne, accuracy*

INTRODUCTION

The increase in greenhouse gases, particularly carbon dioxide, has led to global climate change. The burning of fossil fuels, deforestation, and industrial processes are the primary sources of greenhouse gas emissions. Natural processes, such as the decomposition of organic matter, also release greenhouse gases into the atmosphere. The removal of greenhouse gases from the atmosphere by

Page 28-31 © MANTECH PUBLICATIONS 2023. All Rights Reserved

natural processes, such as photosynthesis and ocean uptake, are called sinks.

Remote sensing has become an essential tool for monitoring greenhouse gases and their sources and sinks on a global scale. Remote sensing involves the collection of data from sensors on satellites, aircraft, and ground-based stations. The data collected provides information on the concentrations of greenhouse gases in the atmosphere and their sources and sinks.

REMOTE SENSING OF CARBON DIOXIDE

Carbon dioxide is the most significant greenhouse gas responsible for climate change. Remote sensing techniques for measuring carbon dioxide include ground-based, airborne, and satellite-based measurements. Ground-based measurements involve the use of Fourier Transform Infrared (FTIR) spectroscopy, which measures the amount of carbon dioxide in the atmosphere by analyzing the absorption of infrared radiation. Airborne measurements involve the use of Lidar (Light Detection and Ranging), which uses laser pulses to measure the concentration of carbon dioxide in the atmosphere. Satellite-based measurements involve the use of passive remote sensing techniques, such as the Orbiting Carbon Observatory-2 (OCO-2) and the Greenhouse Gases Observing Satellite (GOSAT), which use the absorption of sunlight to measure the concentration of carbon dioxide in the atmosphere.

REMOTE SENSING OF METHANE

Methane is the second most significant greenhouse gas responsible for climate change. Methane emissions primarily come from natural sources, such as wetlands and ruminant animals, and human activities, such as fossil fuel extraction and rice cultivation. Remote sensing techniques for measuring methane include ground-based, airborne, and satellite-based measurements. Ground-based measurements involve the use of Fourier Transform Infrared (FTIR) spectroscopy, which measures the amount of methane in the atmosphere by analyzing the absorption of infrared radiation. Airborne measurements involve the use of Lidar (Light Detection and Ranging), which uses laser pulses to measure the concentration of methane in the atmosphere. Satellite-based measurements involve the use of passive remote sensing techniques, such as the Tropospheric Monitoring Instrument (TROPOMI) and the Greenhouse Gases Observing Satellite (GOSAT), which use the absorption of

Page 28-31 © MANTECH PUBLICATIONS 2023. All Rights Reserved

sunlight to measure the concentration of methane in the atmosphere.

REMOTE SENSING OF NITROUS OXIDE:

Nitrous oxide is the third most significant greenhouse gas responsible for climate change. Nitrous oxide emissions primarily come from natural sources, such as soils and oceans, and human activities, such as fertilizer use and industrial processes. Remote sensing techniques for measuring nitrous oxide include ground-based, airborne, and satellite-based measurements. Ground-based measurements involve the use of Fourier Transform Infrared (FTIR) spectroscopy, which measures the amount of nitrous oxide in the atmosphere by analyzing the absorption of infrared radiation.

Airborne measurements involve the use of Lidar (Light Detection and Ranging), which uses laser pulses to measure the concentration of nitrous oxide in the atmosphere. Satellite-based measurements involve the use of passive remote sensing techniques, such as the Tropospheric Monitoring Instrument (TROPOMI) and the Greenhouse Gases Observing Satellite (GOSAT), which use the absorption of sunlight to measure the concentration of nitrous oxide in the atmosphere.

CHALLENGES AND LIMITATIONS

Remote sensing techniques for measuring greenhouse gases and their sources and sinks face several challenges and limitations. The accuracy of measurements depends on various factors, such as the resolution of sensors, atmospheric conditions, and the presence of clouds and aerosols. Additionally, it is challenging to distinguish between natural and anthropogenic sources of greenhouse gases using remote sensing techniques. The spatial and temporal variability of greenhouse gas concentrations also makes it difficult to estimate emissions accurately.

FUTURE PROSPECTS

Remote sensing technology is continually advancing, and new sensors and techniques are being developed to improve the accuracy of greenhouse gas measurements. The European Space Agency's CarbonSat mission, set to launch in 2028, will provide high-resolution measurements of carbon dioxide concentrations. Additionally, the National Aeronautics and Space Administration's (NASA) Geostationary Carbon Cycle Observatory (GeoCARB) mission, set to launch in the mid-2020s, will provide continuous measurements of carbon dioxide, methane, and carbon monoxide

Page 28-31 © MANTECH PUBLICATIONS 2023. All Rights Reserved

over North America. These missions are expected to significantly improve our understanding of greenhouse gases and their sources and sinks.

CONCLUSION

Remote sensing has become an essential tool for monitoring greenhouse gases and their sources and sinks on a global scale. Carbon dioxide, methane, and nitrous oxide are the three most significant greenhouse gases responsible for climate change. Ground-based, airborne, and satellite-based measurements are used to measure greenhouse gases, and their sources and sinks, but these techniques face several challenges and limitations. Advancements in remote sensing technology are expected to improve the accuracy of greenhouse gas measurements in the future. The continuous monitoring of greenhouse gases is crucial for understanding the Earth's climate and developing effective strategies to mitigate climate change.

REFERENCES

1. Ciais, P., Sabine, C., Bala, G., Bopp, L., Brovkin, V., Canadell, J., ... & John, J. (2013). Carbon and other biogeochemical cycles. In Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press.
2. Crisp, D., Atlas, R., Breon, F. M., Brown, L. R., Burrows, J. P., Ciais, P., ... & Rayner, P. (2015). The Orbiting Carbon Observatory-2: First 18 months of science data products. Atmospheric Measurement Techniques, 8(7), 2437-2453.
3. Friedl, M. A., Sulla-Menashe, D., Tan, B., Schneider, A., Ramankutty, N., Sibley, A., & Huang, X. (2010). MODIS Collection 5 global land cover: algorithm refinements and characterization of new datasets. Remote Sensing of Environment, 114(1), 168-182.

Page 28-31 © MANTECH PUBLICATIONS 2023. All Rights Reserved