MANTECH Publications
Journal of Remote Sensing, Environmental Science & Geotechnical Engineering
Volume 5, Issue 2, May-August, 2020

# *Remote Sensing for Monitoring Soil Salinity in Agricultural and Semi-Arid Regions*

***Dr. Ananya Kulkarni***
*Associate Professor*
*Department of Civil and Environmental Engineering*
*Greenfield Institute of Technology, Pune, India*
***Email:*** *ananya.kulkarni@git.edu*

***Rahul Mehta***
*Research Scholar*
*Centre for Geoinformatics and Remote Sensing*
*Sunrise University, Jaipur, India*
***Email:*** *rahul.mehta@sunriseuniv.ac.in*

## *ABSTRACT*

*Soil salinity is a critical environmental and agricultural challenge, especially in arid and semi-arid regions where water scarcity and intensive irrigation accelerate salt accumulation in the soil. Remote sensing technologies, integrated with Geographic Information Systems (GIS), have emerged as powerful tools for mapping, monitoring, and managing salinity over large geographic areas. This paper presents a comprehensive review of remote sensing techniques for soil salinity detection, focusing on spectral indices, microwave sensing, and thermal infrared approaches. It discusses the applicability of satellite platforms such as Landsat, Sentinel, and hyperspectral sensors for salinity mapping, along with the role of GIS in spatial analysis and predictive modeling. The study also examines key factors influencing soil salinity patterns, the integration of field sampling with satellite data, and the development of decision support systems for sustainable agriculture. Case studies from agricultural and semi-arid regions illustrate the*


Page 32-36 © MANTECH PUBLICATIONS 2020. All Rights Reserved

*effectiveness of these methods in guiding salinity mitigation strategies. The paper concludes with recommendations for integrating multi-source remote sensing datasets with machine learning for improved salinity monitoring and management.*

***Keywords:*** *Soil salinity, Remote sensing, Semi-arid regions, Spectral indices, GIS, Hyperspectral imaging, Precision agriculture*

## INTRODUCTION

Soil salinity poses a significant constraint to agricultural productivity worldwide, particularly in arid and semi-arid regions where irrigation is essential for crop cultivation. Excessive salt concentration in soil not only hampers plant growth but also degrades soil structure, leading to reduced infiltration, increased erosion, and long-term desertification. Traditionally, salinity assessment has relied on field surveys and laboratory analysis of soil samples. While accurate, these methods are time-consuming, labor-intensive, and limited in spatial coverage.

Remote sensing offers a cost-effective and efficient alternative by providing repetitive, synoptic, and multi-spectral data that can be processed to detect salinity patterns at various spatial scales. With advancements in satellite technology, high-resolution multispectral and hyperspectral imagery, microwave data, and thermal infrared measurements are now widely accessible for environmental monitoring. When integrated with GIS, these datasets enable spatial modeling, change detection, and the creation of salinity risk maps to guide land management decisions.

## SOIL SALINITY PROBLEM IN AGRICULTURAL AND SEMI-ARID REGIONS

Salinity in agricultural soils results from natural processes such as weathering of parent rocks, capillary rise of saline groundwater, and marine influence, as well as human-induced factors like over-irrigation, poor drainage, and use of saline water for irrigation. Semi-arid regions, characterized by low rainfall and high evapotranspiration rates, are especially vulnerable to salinization due to insufficient leaching of salts from the root zone.

Impacts of soil salinity include:

- Reduced crop yields and crop failure
- Altered soil microbial activity
- Soil dispersion and crusting
- Decline in soil organic matter


Page 32-36 © MANTECH PUBLICATIONS 2020. All Rights Reserved

## REMOTE SENSING TECHNIQUES FOR SALINITY DETECTION

### Spectral Characteristics of Saline Soils

Saline soils display unique spectral signatures in the visible, near-infrared (NIR), and short-wave infrared (SWIR) regions. The presence of salts often increases reflectance in the visible range and reduces it in the NIR region due to surface crusting and salt efflorescence.

### Multispectral Remote Sensing

Multispectral data from platforms such as Landsat (TM, ETM+, OLI) and Sentinel-2 provide medium-resolution imagery useful for regional salinity mapping. Vegetation indices like NDVI and salinity-specific indices such as SI (Salinity Index) and NDSI (Normalized Difference Salinity Index) are widely applied.

### Hyperspectral Imaging

Hyperspectral sensors capture hundreds of narrow spectral bands, enabling detailed identification of soil mineral composition and moisture status. Instruments like Hyperion and upcoming missions such as EnMAP provide high spectral resolution data suitable for mapping varying salinity levels.

### Microwave and Thermal Infrared Sensing

Active microwave sensors such as Synthetic Aperture Radar (SAR) are effective under all-weather conditions, detecting changes in soil surface roughness and moisture, which correlate with salinity. Thermal infrared data can be used to infer soil moisture stress and evaporation rates, indirectly indicating salinity severity.

## ROLE OF GIS IN SALINITY MONITORING

GIS plays an essential role in processing, analyzing, and visualizing salinity data. By integrating satellite imagery with ancillary datasets such as soil maps, topography, and climate data, GIS enables spatial modeling of salinity risk zones.

Key GIS applications include:

- Interpolation of field-measured salinity values
- Trend analysis of salinity over time
- Overlay analysis to determine salinity-crop yield relationships
- Development of decision support systems for farmers


Page 32-36 © MANTECH PUBLICATIONS 2020. All Rights Reserved

## CASE STUDIES

### Case Study 1: Punjab, India

Using Landsat-8 OLI imagery and field EC measurements, researchers developed a salinity index map for irrigated fields in Punjab. Results indicated that over 15% of the area showed moderate to high salinity, correlating strongly with poor drainage and over-irrigation practices.

### Case Study 2: Murray-Darling Basin, Australia

Hyperspectral data were used to map saline discharge zones in the basin. The integration of remote sensing with groundwater models improved the prediction of future salinity hotspots and guided irrigation scheduling.

## TABLE 1: Common Remote Sensing Indices for Soil Salinity Detection

| Index | Formula | Description |
|---|---|---|
| Salinity Index (SI) | (Red × NIR) | Combines visible and NIR reflectance to detect salt crusts |
| Normalized Difference Salinity Index (NDSI) | (Red − SWIR) / (Red + SWIR) | Detects high salt content in soils |
| Vegetation Salinity Index (VSI) | (NIR / Green) | Indicates vegetation stress due to salinity |

## ADVANTAGES AND LIMITATIONS

**Advantages**:

- Large area coverage
- Repetitive monitoring
- Cost-effective compared to ground surveys
- Multi-temporal analysis possible

**Limitations**:

- Influence of soil moisture and vegetation cover on reflectance
- Atmospheric effects on spectral data
- Requirement of field calibration and validation


Page 32-36 © MANTECH PUBLICATIONS 2020. All Rights Reserved

## FUTURE PROSPECTS

The integration of artificial intelligence and machine learning with remote sensing and GIS is expected to revolutionize salinity mapping. Deep learning algorithms can analyze multi-source datasets to improve classification accuracy and predict salinity dynamics under climate change scenarios. The increasing availability of high-resolution imagery from CubeSats and commercial satellites will enable near real-time salinity monitoring, aiding precision agriculture.

## CONCLUSION

Remote sensing combined with GIS offers a powerful framework for monitoring soil salinity in agricultural and semi-arid regions. The use of spectral indices, hyperspectral imaging, and microwave sensing provides multiple approaches for detecting salinity patterns. GIS facilitates the integration of spatial datasets, enabling comprehensive analysis and targeted mitigation strategies. The adoption of these technologies, coupled with ground truthing and advanced analytics, will be essential for sustainable agricultural development in salinity-prone areas.

## REFERENCES

[1] Metternicht, G.I., Zinck, J.A., "Remote sensing of soil salinity: potentials and constraints," *Remote Sensing of Environment*, vol. 85, no. 1, pp. 1–20, 2003.
[2] Abbas, A., Khan, S., Hussain, N., "Characterizing soil salinity in irrigated areas using remote sensing," *Journal of Environmental Management*, vol. 88, pp. 130–138, 2008.
[3] Allbed, A., Kumar, L., "Soil salinity mapping and monitoring in arid and semi-arid regions using remote sensing technology: A review," *Advances in Remote Sensing*, vol. 2, pp. 373–385, 2013.
[4] Bannari, A., et al., "Detection of soil salinity using airborne hyperspectral data," *Remote Sensing of Environment*, vol. 96, pp. 489–509, 2005.
[5] Dwivedi, R.S., "Study of salt-affected soils using remote sensing techniques," *International Journal of Remote Sensing*, vol. 19, pp. 195–209, 1998.
[6] Weng, Y., Gong, P., Zhu, Z., "A remote sensing–GIS evaluation of soil salinity in an arid region of China," *Environmental Monitoring and Assessment*, vol. 91, pp. 335–349, 2004.
[7] Bouaziz, M., et al., "Monitoring soil salinity in irrigated agriculture using multi-temporal remote sensing data," *Agricultural Water Management*, vol. 97, pp. 139–147, 2010.


Page 32-36 © MANTECH PUBLICATIONS 2020. All Rights Reserved