MANTECH Publications
Journal of Remote Sensing, Environmental Science & Geotechnical Engineering
Volume 1, Issue 3, September-December, 2016

Climate Change Impact Assessment Using Multi-Temporal Remote Sensing Data

Dr. Ananya Mehta
Associate Professor
Department of Civil Engineering
Greenfield Institute of Technology, Pune, India.
Email:*ananya.mehta@gitp.ac.in*

Dr. Raghav Sharma
Assistant Professor
Department of Environmental Science
Horizon College of Engineering, Jaipur, India.
Email:*raghav.sharma@hce.ac.in*

Abstract

Climate change presents unprecedented challenges for environmental sustainability, ecosystem stability, and socio-economic development. Multi-temporal remote sensing data offers a robust approach to assessing the impacts of climate change through the continuous monitoring of land, water, and atmospheric variables over time. By integrating satellite-based observations with geospatial analysis, scientists can identify long-term trends in temperature variation, vegetation health, glacier retreat, sea-level rise, and urban heat dynamics. This paper examines the application of multi-temporal remote sensing in climate change impact assessment, discussing key datasets, analytical methods, case studies, and the integration of remote sensing with predictive climate models. The analysis highlights how spatially explicit and time-series data can support early warning systems, adaptive resource management, and policy development to mitigate climate-related risks.

Page 82-86 © MANTECH PUBLIATIONS 2016. All Rights Reserved

Keywords: *Multi-temporal remote sensing, climate change, impact assessment, geospatial analysis, land surface temperature, vegetation dynamics, environmental monitoring.*

INTRODUCTION

Climate change is one of the most significant global environmental threats, affecting biodiversity, agricultural productivity, water resources, and human settlements. The ability to detect, quantify, and analyze these impacts requires continuous monitoring of the Earth's surface. Remote sensing has revolutionized environmental monitoring by providing consistent, repetitive, and large-scale observations.

Multi-temporal remote sensing refers to the use of satellite imagery acquired over different time periods to detect and analyze changes in the Earth's systems. Such data allow researchers to track seasonal patterns, long-term climate shifts, and extreme weather events. Unlike ground-based monitoring, satellite-based approaches offer cost-effective, spatially comprehensive datasets for regional to global assessments.

This paper focuses on the methods, applications, and case studies of using multi-temporal remote sensing for climate change impact assessment.

REMOTE SENSING FOR CLIMATE MONITORING

Remote sensing systems capture electromagnetic radiation reflected or emitted from the Earth's surface. Optical, thermal, and microwave sensors provide different layers of information that can be used to study climate-related phenomena.

Optical sensors (e.g., Landsat, Sentinel-2) monitor vegetation and land cover. **Thermal sensors** (e.g., MODIS, Landsat TIRS) assess land surface temperature (LST). **Microwave sensors** (e.g., Sentinel-1 SAR) are useful for detecting soil moisture and ice sheet changes.

MULTI-TEMPORAL DATASETS

The following satellite missions provide valuable multi-temporal datasets for climate change studies:

Page 82-86 © MANTECH PUBLIATIONS 2016. All Rights Reserved

Satellite	Key Parameters	Temporal Resolution	Application
Landsat Series	Land cover, LST, NDVI	16 days	Vegetation change, urban expansion
MODIS	LST, vegetation indices, snow cover	Daily	Seasonal & annual trend analysis
Sentinel-2	Land cover, vegetation health	5 days	Crop monitoring, forest dynamics
Sentinel-1 SAR	Soil moisture, ice monitoring	6–12 days	Flood mapping, glacier retreat

Table 1: Commonly used multi-temporal remote sensing datasets for climate monitoring.

METHODS FOR IMPACT ASSESSMENT

The assessment of climate change impacts using multi-temporal remote sensing involves several analytical methods:

Change Detection Analysis

Detecting differences in satellite images over time to identify trends in vegetation, water bodies, or glaciers.

Time-Series Analysis

Examining continuous datasets to identify periodic or long-term variations.

Anomaly Detection

Recognizing unusual deviations from historical averages, such as extreme droughts or sudden vegetation loss.

Integration with Climate Models

Coupling remote sensing observations with predictive models for improved climate projections.

APPLICATIONS IN CLIMATE CHANGE STUDIES

1. Glacier Retreat Monitoring

Page 82-86 © MANTECH PUBLIATIONS 2016. All Rights Reserved

High-resolution optical and SAR data reveal spatial patterns of glacier loss in Himalayan regions, indicating reduced freshwater availability.

2. Vegetation Health Assessment

NDVI time-series data help detect shifts in vegetation productivity due to temperature and precipitation changes.

3. Coastal Change Detection

Multi-temporal Landsat and Sentinel imagery track shoreline retreat and coastal flooding associated with sea-level rise.

4. Urban Heat Island Studies

Thermal imagery quantifies temperature anomalies in urban areas and their growth trends.

CASE STUDY: VEGETATION RESPONSE TO CLIMATE VARIABILITY

In a study across central India, Landsat-based NDVI time-series (2000–2023) showed declining vegetation health in semi-arid regions due to reduced monsoon rainfall and increased land surface temperatures.

ADVANTAGES AND LIMITATIONS

Advantages

- Synoptic and repetitive coverage.
- Cost-effective and globally accessible datasets.
- Ability to integrate with GIS and climate models.

Limitations

- Cloud cover interference in optical imagery.
- Need for high technical expertise for data processing.
- Limited temporal resolution in some satellites.

FUTURE DIRECTIONS

Advances in remote sensing technology, including hyperspectral imaging and constellations of small satellites, will enable higher-frequency and higher-resolution monitoring. Machine

Page 82-86 © MANTECH PUBLIATIONS 2016. All Rights Reserved

learning and AI will play a key role in automating change detection and predicting future climate impacts.

CONCLUSION

Multi-temporal remote sensing has become a cornerstone for climate change impact assessment, offering objective, large-scale, and repeatable observations. By integrating data from multiple satellites and applying advanced analytical techniques, scientists can detect environmental changes, assess vulnerability, and guide climate adaptation strategies. As climate change accelerates, multi-temporal remote sensing will be indispensable for sustainable resource management and informed policymaking.

REFERENCES

1. Justice, C. O., et al. "The Moderate Resolution Imaging Spectroradiometer (MODIS): Land Remote Sensing for Global Change Research." *IEEE Transactions on Geoscience and Remote Sensing*, vol. 36, no. 4, 1998, pp. 1228–1249.
2. Wulder, M. A., et al. "Landsat Continuity: Contributions of the Landsat Program to Earth Observation Science and Applications." *Remote Sensing of Environment*, vol. 122, 2012, pp. 1–6.
3. Singh, R. K., and D. S. Roy. "Remote Sensing of Climate Change Impacts in India." *International Journal of Applied Earth Observation and Geoinformation*, vol. 45, 2016, pp. 45–54.
4. Hansen, M. C., et al. "High-Resolution Global Maps of 21st-Century Forest Cover Change." *Science*, vol. 342, 2013, pp. 850–853.
5. Fensholt, R., et al. "Greenness in Semi-Arid Areas across the Globe 1981–2007—An Earth Observing Satellite Based Analysis of Trends and Drivers." *Remote Sensing of Environment*, vol. 114, no. 7, 2010, pp. 1286–1296.
6. Small, C., et al. "Urban Remote Sensing: Overview and Challenges." *Remote Sensing of Environment*, vol. 86, no. 3, 2003, pp. 273–283.
7. Liang, S. "Quantitative Remote Sensing of Land Surfaces." John Wiley & Sons, 2004.

Page 82-86 © MANTECH PUBLIATIONS 2016. All Rights Reserved